**Australian Government** 



Australian Fisheries Management Authority



## Commonwealth Small Pelagic Fishery: Fishery Assessment Report 2019–2021

Report to the Australian Fisheries Management Authority

G. L. Grammer, T. M. Ward and L. M. Durante

RR2019/0803 June 2022

## Contents

Acknow	Acknowledgements				
Executiv	Executive Summary5				
1 Gen	1 General Introduction				
1.1	Overview	8			
1.2	Description of the Commonwealth Small Pelagic Fishery	8			
1.3	Management of the Fishery	9			
1.4	Previous Assessments	11			
1.5	Aims and Objectives	12			
2 Jack	mackerel (Trachurus declivis)	13			
2.1	Introduction	13			
2.2	Methods	19			
2.3	Results	21			
2.4	Summary and Conclusions	33			
3 Blue	e Mackerel (Scomber australasicus)	35			
3.1	Introduction	35			
3.2	Methods	40			
3.3	Results	42			
3.4	Summary and Conclusions	54			
4 Red	bait (Emmelichthys nitidus)	56			
4.1	Introduction	56			
4.2	Methods	61			
4.3	Results	63			
4.4	Summary and Conclusions	75			
5 Aus	tralian Sardine (Sardinops sagax)	76			
5.1	Introduction	76			
5.2	Methods	81			
5.3	Results	82			
5.4	Summary and Conclusions	89			
6 Gen	eral Summary and Conclusions				
Referenc	ces	91			

This publication may be cited as:

Grammer, G. L., Ward, T.M. and Durante, L.M. (2022). Commonwealth Small Pelagic Fishery: Fishery Assessment Report 2019-2021. Report to the Australian Fisheries Management Authority. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2010/000270-11. SARDI Research Report Series No. 1133. 101pp.

#### AUTHOR AFFILIATIONS

<u>Gretchen L. Grammer and Leonardo M. Durante:</u> South Australian Research and Development Institute (Aquatic Sciences), 2 Hamra Ave., West Beach, SA 5024, Australia

<u>Timothy M. Ward:</u> Institute for Marine and Antarctic Studies, University of Tasmania, Private Bag 49, Hobart, Tasmania 7001, Australia

#### DISCLAIMER

The authors warrant that they have taken all reasonable care in producing this report. The report has been through the SARDI internal review process, and has been formally approved for release by the Research Director, Aquatic Sciences. Although all reasonable efforts have been made to ensure quality, SARDI does not warrant that the information in this report is free from errors or omissions. SARDI and its employees do not warrant or make any representation regarding the use, or results of the use, of the information contained herein as regards to its correctness, accuracy, reliability and currency or otherwise. SARDI and its employees expressly disclaim all liability or responsibility to any person using the information or advice. Use of the information and data contained in this report is at the user's sole risk. If users rely on the information they are responsible for ensuring by independent verification its accuracy, currency or completeness. The SARDI Report Series is an Administrative Report Series which has not been reviewed outside the department and is not considered peer-reviewed literature. Material presented in these Administrative Reports may later be published in formal peer-reviewed scientific literature.

#### © 2022 SARDI & AFMA

This work is copyright. Apart from any use as permitted under the *Copyright Act* 1968 (Cth), no part may be reproduced by any process, electronic or otherwise, without the specific written permission of the copyright owner. Neither may information be stored electronically in any form whatsoever without such permission.

Author(s):G.L. Grammer, T.M. Ward and L. M. DuranteReviewer(s):L. McLeay and F. Bailleul (SARDI) and H. Johnston (AFMA)Approved by:S. Mayfield<br/>Science Leader – FisheriesSigned:Mayfield<br/>Science Leader – FisheriesDate:16 June 2022Distribution:AFMA, SARDI Aquatic Sciences, Parliamentary Library, State Library and National<br/>LibraryCirculation:OFFICIALALL ENQUIRIES

South Australian Research and Development Institute - SARDI Aquatic Sciences 2 Hamra Avenue West Beach SA 5024 PO Box 120 Henley Beach SA 5022 **P:** (08) 8207 5400 **F:** (08) 8207 5415 **E:** <u>pirsa.sardiaquatics@sa.gov.au</u> **W:** <u>http://www.pir.sa.gov.au/research</u>

# Acknowledgements

This fishery assessment report was funded by the Australian Fisheries Management Authority (AFMA). Data presented in this report were provided by: Mr Max Bayly (Australian Fisheries Management Authority); Ms Denise Garcia (Department of Primary Industries, Parks, Water and Environment Tasmania); Dr John Stewart (New South Wales Department of Primary Industries); Ms Monique Nelis (Victorian Department of Economic Development, Jobs, Transport and Resources); and Mr Angelo Tsolos (SARDI Aquatic Sciences). Catch samples were processed by Mr Alex Ivey, Mr Andrew Hutchinson and Ms Sharon Drabsch. This report was formally reviewed by Dr Lachlan McLeay and Dr Fred Bailleul and approved for release by Dr Stephen Mayfield (SARDI Aquatic Sciences).

# **Executive Summary**

This report uses fishery-dependent and fishery-independent data to assess the status of target species in the East and West sub-areas of the Commonwealth Small Pelagic Fishery (SPF) from 2018/19 to 2020/21. This report updates the fishery assessment report of Ward and Grammer (2019) and the status summary reports of Ward and Grammer (2021a, 2021b).

The target species in the SPF are Jack Mackerel (*Trachurus declivis*), Blue Mackerel (*Scomber australasicus*), Redbait (*Emmelichthys nitidus*) and, in the Australian Sardine sub-area of the East sub-area, Australian Sardine (*Sardinops sagax*). The primary fishing methods are purse-seining and mid-water trawling. The assessment uses commercial catch and effort data for the SPF up to 30 April 2021 and biological information (size and age structures, reproduction). Data from State-managed commercial fisheries are used to estimate the total catch of each species in each sub-area.

The SPF Harvest Strategy 2008 (last revised April 2017) specifies that the primary technique for assessing the status of SPF species is the Daily Egg Production Method (DEPM). To retain a species in a sub-area at Tier 1, where exploitation rates are highest, the DEPM must be applied every 5 years. Between applications of the DEPM, fishery-dependent data are analysed to identify variations in fishing patterns or catches that may be indicative of changes in stock status and/or localised depletion.

Between 2010/11 and 2014/15, catches in the SPF of all species in both sub-areas were low (<200 t per annum). Effort and catch in the SPF increased during 2015/16 and 2016/17, when a mid-water factory trawler operated in offshore waters of both sub-areas. Since 2017/18, smaller mid-water trawlers without onboard processing facilities began operating in inshore waters of the East sub-area. Negligible fishing was undertaken by SPF vessels in the West sub-area from 2017/18 to 2020/21.

## Jack Mackerel

The estimate of spawning biomass for Jack Mackerel East in 2019 was 156,292 t (95% CI = 49,120–263,496) (Ward et al. 2020a). The total catch of Jack Mackerel East in 2020/21 was 5,894 t. The total catch in 2019/20 was 7,852 t, which was the highest catch in more than 20 years (i.e. since 1997/98). The SPF is the main fishery for Jack Mackerel in the East sub-area. During 2020/21, the modal size of Jack Mackerel East taken in mid-water trawls (~190 mm FL) was below the approximate mean size at 50% maturity (~230 mm FL). Recent increases in catches of Jack Mackerel East reflect increases in fishing effort. The total catch of Jack Mackerel East was 4% of the 2019 estimate of spawning biomass and below the Tier 1 exploitation rate for this stock of 12%. The total catch by the SPF in 2020/21 was 32% of the available TAC. On the basis of the information provided here, Jack Mackerel East is classified as **sustainable**.

The spawning biomass of Jack Mackerel West in 2016/17 was estimated to be at least 31,000 t (Ward et al. 2018). There has been minimal to no catch of Jack Mackerel West by the SPF since 2017/18. The highest recent catches were 686 t in 2016/17 and 634 t in 2015/16. The SPF has been the main fishery for Jack Mackerel in the West sub-area. Recent annual catches of Jack Mackerel West have been <1% of the 2016/17 estimate of spawning biomass and below the Tier 1 exploitation rate for this stock of 12%. The total catch in 2020/21 by the SPF was 0% of the available TAC. On the basis of the information provided here, Jack Mackerel West is classified as **sustainable**.

## **Blue Mackerel**

The spawning biomass of Blue Mackerel East in 2019 was ~80,000 t (Ward et al. 2021a). The total catch of Blue Mackerel East in 2020/21 was 6,202 t, which is the highest on record. The second highest catch was 6,124 in 2019/20. The main fisheries for Blue Mackerel in the East sub-area are the SPF (93% in 2020/21) and the New South Wales (NSW) Ocean Hauling Fishery (7% in 2020/21). During 2019/20, the modal size of Blue Mackerel East from mid-water trawls (250 mm FL) was similar to the mean size at 50% maturity (~260 mm FL). Recent increases in catches of Blue Mackerel East reflect increases in fishing effort. The total catch of Blue Mackerel East in 2020/21 was 8% of the 2019 estimate of spawning biomass and below the Tier 1 exploitation rate for this stock of 15%. The total catch by the SPF in 2020/21 was 52% of the available TAC. On the basis of the information provided here, Blue Mackerel East is classified as **sustainable**.

The spawning biomass for Blue Mackerel West in 2005 was estimated to be 56,228 t (Ward et al. 2009). There has been minimal to no catch of Blue Mackerel West by the SPF since 2017/18. The highest recent catches were 766 t in 2016/17 and 980 t in 2015/16. The SPF has been the main fishery for Blue Mackerel in the West sub-area. Recent total annual catches of Blue Mackerel West have been <1% of the estimated spawning biomass for 2005 and below the Tier 3 exploitation rate for this stock of 3.75%. The total catch in 2020/21 was 0% of the available TAC. On the basis of the information provided here, Blue Mackerel West is classified as **sustainable**.

## Redbait

The estimate of spawning biomass for Redbait East in 2020 was 52,629 t (95% CI = 13,937-91,321) (Grammer et al. 2022). The total catch of Redbait East in 2020/21 was 2,011 t. The total catch in 2019/20 was 2,457 t, which is the highest catch since 2006/07, but well below the peak of 7,733 t in 2003/04. The SPF is the main fishery for Redbait in the East sub-area. During 2020/21, the modal size of Redbait East from mid-water trawls (180 mm FL) was above the mean size at 50% maturity (~150 mm FL). The recent increase in catches of Redbait East reflect increases in fishing effort. The total catch of Redbait East in 2020/21 by the SPF was 4% of the 2020 estimate of spawning biomass and below the Tier 2 exploitation rate for this stock of 5%. The total catch by the SPF in

2020/21 was 59% of the available TAC. On the basis of the information provided here, Redbait East is classified as **sustainable**.

The estimate of spawning biomass for Redbait West in 2017 was 66,767 t (CI = 28,797-190,392) (Ward et al. 2019). There has been minimal to no catch of Redbait West by the SPF since 2017/18. The highest recent catches were 1,140 t in 2016/17 and 1,157 t in 2015/16. The SPF has been the main fishery for Redbait in the West sub-area. Recent catches of Redbait West have been <1% of the 2017 estimate of spawning biomass and below the Tier 1 exploitation rate for this stock of 10%. The total catch in 2020/21 by the SPF was 0% of the available TAC. On the basis of the information provided here, Redbait West is classified as **sustainable**.

## **Australian Sardine**

The estimate of spawning biomass of Australian Sardine in the Sardine sub-area in 2019 was 42,724 t (95% CI = 15,487–69,962 t) (Ward et al. 2021a). The total catch of Sardine in the Sardine sub-area in 2020/21 was 717 t and 733 t in 2019/20, which were the highest annual catch since 2010/11 (1,574 t). The main fisheries for Sardine in the Sardine sub-area are the SPF (12% in 2020/21) and the NSW Ocean Hauling Fishery (88% in 2020/21). Catches of Sardine from the Sardine sub-area have mainly comprised fish at or above the mean size at 50% maturity of ~150 mm FL. The total catch of Sardine in the Sardine sub-area was <2% of the 2020 estimate of spawning biomass and below the Tier 1 exploitation rate for this stock of 20%. The total catch by the SPF in 2020/21 was <1% of the available TAC. On the basis of the information provided here, Sardine in the Sardine sub-area is classified as **sustainable**.

## Summary

All SPF stocks are classified as sustainable. DEPM surveys have been conducted for all stocks.

Redbait East returned to Tier 1 in 2022/23 following a DEPM survey in October 2020. All SPF stocks are at Tier 1 except Blue Mackerel West. The DEPM was last applied to Blue Mackerel West in 2005. Estimates of spawning biomass obtained from DEPM surveys could be used to inform the establishment of target (e.g. B<sub>50</sub>) and limit reference points (e.g. B<sub>20</sub>) for each stock.

**Keywords:** Commonwealth SPF, Jack Mackerel, Blue Mackerel, Redbait, Sardine, purse seine, mid-water trawl, AFMA.

# **1** General Introduction

## 1.1 Overview

This assessment of the Commonwealth Small Pelagic Fishery (SPF) builds on annual reports published since 2010 (Ward et al. 2011b, 2012, 2013, 2014c, 2015c; Ward and Grammer 2016, 2017, 2018, 2019, 2021a, 2021b). This report provides a synopsis of information available and current status of SPF quota species, namely Jack Mackerel (*Trachurus declivis*), Blue Mackerel (*Scomber australasicus*), Redbait (*Emmelichthys nitidus*) and Australian Sardine (*Sardinops sagax*). The assessment uses commercial catch and effort data up to 30 April 2021 and available biological information (size and age structures, reproduction). The assessments are underpinned by outputs from several Management Strategy Evaluations (MSEs) and estimates of spawning biomass obtained from fishery-independent surveys. This report satisfies the requirements of the SPF Harvest Strategy (Harvest Strategy) for assessment of stocks at Tiers 1 and 2 (see Section 1.3.1; AFMA 2008, revised in 2017).

## 1.2 Description of the Commonwealth Small Pelagic Fishery

The SPF is a purse seine and mid-water trawl fishery that operates in Commonwealth waters (3 to 200 nm) from southern Queensland to south-western Western Australia, including Tasmania (Figure 1-1). The fishery is divided into two sub-areas (East and West) by a line through longitude 146°30'E (Figure 1-1, AFMA 2009). There is also a designated Australian Sardine sub-area within the East sub-area that extends from southern Queensland to southern New South Wales (NSW) (Figure 1-1; referred to here as the 'Sardine East sub-area'). The East and West sub-areas are further divided into seven sub-zones and catch grids (Figure 1-2).

The three main target species of the SPF are Jack Mackerel, Blue Mackerel and Redbait. Australian Sardine is a target species in the Sardine East sub-area. These species are also targeted by recreational fishers in some States (Henry and Lyle 2003) and by Statemanaged commercial fisheries. Species caught in the SPF go into the domestic market as fishmeal, bait and for human consumption (Patterson et al. 2021).

Small quantities of SPF species are caught in other Commonwealth fisheries, primarily the Southern and Eastern Scalefish and Shark Fishery, Western Tuna and Billfish Fishery and the Eastern Tuna and Billfish Fishery (Moore and Skirtun 2012). Combined catches of SPF quota species in these fisheries have not exceed 40 t per year since their inception in 2002 and are not included in this assessment.

The SPF has changed dramatically since large scale fishing operations targeting Jack Mackerel began in the mid-1980s. Between the late 1980s and prior to the SPF Management Plan being implemented in 2009 (AFMA 2009), the Tasmanian component of the fishery (north-eastern Tasmania to central western Tasmania) was managed using a combination of input and output controls, including a total allowable catch (TAC). A combined species TAC for the Tasmanian component of the fishery was set at 42,000 t in 1988/89 and was based on the highest annual catch from the purse seine fishery (Jordan et al. 1992; Pullen 1994). The TAC was decreased to 34,000 t in 2002/03 with the renewed interest in small pelagic species and the commencement of mid-water trawl operations. Despite catches not approaching this level, this TAC was applied in subsequent fishing seasons up until 2008/09 when the SPF was split into East and West sub-areas (SPF Management Plan; AFMA 2009). TACs specific to each species and sub-area were applied as outlined by the Harvest Strategy (AFMA 2008).

Effort and catch in the SPF have fluctuated over time, driven by a combination of social, economic, biological and ecological factors. A detailed history of the SPF is described in Moore and Skirtun (2012). Since 2014, effort has increased in the SPF. Catches increased in 2014/15 to 2015/16 when a factory trawler operated in both sub-areas of the SPF (Ward and Grammer 2019), and again in 2016/17 when a new fishing operation was established in the East sub-area off southern NSW. Since 2017/18, smaller mid-water trawlers have taken regular catches in the East sub-area of the SPF (Ward and Grammer 2021a).

### 1.3 Management of the Fishery

The SPF is managed by the Australian Fisheries Management Authority (AFMA) under the SPF Management Plan (AFMA 2009) using a combination of input and output controls that include limited entry, zoning, mesh size restrictions and TAC limits for target species (hereafter referred to as quota species) within each sub-area.



Figure 1-1 Management sub-areas of the Small Pelagic Fishery.



Figure 1-2. Sub-zones, catch grids and areas closed to mid-water trawling within management sub-areas of the Small Pelagic Fishery.

#### 1.3.1 Harvest Strategy

The Harvest Strategy for the SPF, established in 2008, was last revised in 2017 (AFMA 2008, revised April 2017). The Harvest Strategy is a three-tiered system used by the SPF Resource Assessment Group (SPFRAG; previously the SPF Scientific Panel, SPFSP) to develop advice on the Recommended Biological Catches (RBCs) for stocks (East and West) for each quota species. Stocks are allocated to a tier based upon the level of knowledge about stock size (spawning biomass), with Tier 1 representing the highest level of available information and Tier 3 the lowest (Moore and Skirtun 2012). Corresponding individual transferable quotas (ITQs) are established; Tier 1 stocks have the largest quota (by weight), and Tier 3 the smallest (Tracey et al. 2013). The tiered system was introduced to ensure that heavy exploitation only occurs in stocks where there is a high level of confidence that such exploitation can be sustained (Moore and Skirtun 2012). TACs for each quota species are determined by subtracting other sources of mortality (i.e. catches taken in other Commonwealth and State fisheries) from the corresponding RBCs.

A brief description of each tier is provided below.

<u>Tier 1:</u> The maximum exploitation rates for Tier 1 species in each sub-area are 10% for Redbait, 12% for Jack Mackerel, 15% for Blue Mackerel and 20% for Australian Sardine (East sub-area only). RBCs are set by applying exploitation rates up to these levels based on the median spawning biomass estimated using the Daily Egg Production Method (DEPM). Species remain at Tier 1 for five seasons after a DEPM survey is completed.

<u>Tier 2:</u> The maximum exploitation rates for Tier 2 species are half the level specified at Tier 1. Redbait and Jack Mackerel can remain at Tier 2 for up to 10 seasons. Blue Mackerel and Australian Sardine can remain at Tier 2 for up to 5 seasons.

<u>Tier 3:</u> The maximum exploitation rates for Tier 3 species are half the level specified at Tier 2 when a biomass estimate has been previously based on a DEPM survey, i.e. 2.5% for Redbait, 3% for Jack Mackerel, 3.75% for Blue Mackerel and 5% Australian Sardine (East sub-area only). For stocks with no previous DEPM survey, the exploitation rates may not exceed a quarter of the mean biomass estimate derived from the Atlantis-SPF model, i.e. West sub-area: 1.25% Redbait and 1.5% Jack Mackerel. A stock can remain at Tier 3 indefinitely. The Atlantis-SPF model is a variant of the Atlantis-SE ecosystem model (see section 1.4.2; Smith et al. 2015).

### 1.4 Previous Assessments

#### 1.4.1 DEPM

DEPM surveys have been conducted for Blue Mackerel East and West (Ward and Rogers 2007, Ward et al. 2009, 2015b, 2021a), Australian Sardine East (Ward et al. 2007, 2015a, 2015b, 2021a), Redbait East and West (Neira and Lyle 2011, Ward et al. 2019, Grammer et al. 2022), and Jack Mackerel East and West (Neira 2011, Ward et al. 2015a, 2016, 2018, 2020a).

#### 1.4.2 Management Strategy Evaluations (MSEs)

Management Strategy Evaluations (MSEs) have been conducted by the Australian Bureau of Agricultural and Resource Economics and Sciences (ABARES) (Giannini et al. 2010) and Commonwealth Scientific and Industrial Research Organisation (CSIRO) (Smith et al. 2015). The 2015 MSE included a formal review of the Harvest Strategy for the SPF; these results are discussed in this assessment report.

#### Giannini et al. (2010)

In 2010, an MSE model was used to test the settings (i.e. exploitation rates) in the Harvest Strategy for each stock (Giannini et al. 2010). In most scenarios, the 30 year simulation period used in the MSE was sufficient for each stock to reach equilibrium, and generally this was well above 20% of virgin biomass (B<sub>20</sub>). Sensitivities of the model to the various input parameters were also tested. The model was most sensitive to the assumed stock-

recruitment relationship and natural mortality. The model was re-examined in 2011 to address concerns about values used for number of recruits.

#### Smith et al. (2015)

In 2015, an MSE was undertaken using ecosystem and population models to evaluate and provide advice on the reference points (e.g. biomass depletion levels) and settings (e.g. exploitation rates) for the SPF target species (Smith et al. 2015). A new variant of the Atlantis ecosystem model (Atlantis-SPF) indicated that SPF species are not keystone species within the ecosystem, and population modelling suggested that conventional single species targets and limits (e.g. the defaults under the Harvest Strategy) are appropriate (Smith et al. 2015). Based on results from the ecosystem model and default settings in the Commonwealth Fisheries Harvest Strategy Policy, Smith et al. (2015) evaluated constant Tier 1 exploitation rates for each species that achieved a median depletion to 50% of unfished levels ( $B_{50}$ ) while maintaining a <10% chance of falling below 20% of unfished levels (B<sub>20</sub>). The base case model assumed DEPM surveys every 5 years. Evaluation of the Tier 2 exploitation rate (50% of Tier 1) assumed that it would only be applied after 5 years of exploitation at Tier 1 and no further DEPM surveys would occur. Smith et al. (2015) also produced Atlantis-SPF biomass estimates that have been used in Tier 3 of the Harvest Strategy when DEPM spawning biomass estimates were not available.

Smith et al. (2015) suggested the previous Tier 1 harvest rate of 15% was too high for Jack Mackerel and Redbait and too low for Blue Mackerel and Australian Sardine. Smith et al. (2015) recommended: 1) Tier 1 harvest rates be applied for not more than 5 years after a DEPM survey; 2) Tier 2 harvest rates be 50% of Tier 1 rates; and 3) Tier 2 harvest rates should not be applied for more than 5 years for Blue Mackerel and Australian Sardine or 10 years for Jack Mackerel and Redbait.

## 1.5 Aims and Objectives

This report uses fishery-dependent and fishery-independent data to assess the status of target species of the SPF. The status of Jack Mackerel, Blue Mackerel, Redbait are assessed in the East and West sub-areas. Australian Sardine is assessed in the Sardine sub-area of the East sub-area. This report satisfies the requirements of the Harvest Strategy (AFMA 2008, revised April 2017).

# 2 Jack mackerel (*Trachurus declivis*)

## 2.1 Introduction

#### 2.1.1 Background to Fishery

A large purse seine fishery for small pelagic fishes was developed off Tasmania in the mid-1980s. The majority of the catch was Jack Mackerel (*Trachurus declivis*), with relatively small quantities of Redbait (*Emmelichthys nitidus*) and Blue Mackerel (*Scomber australasicus*) taken as by-product. The fishery became the largest in Australia by weight, with catches of Jack Mackerel peaking at 39,747 t in 1986/87 (Kailola et al. 1993, Pullen 1994). In 1988/89, the Jack Mackerel catch fell to 8,150 t (Kailola et al. 1993, Pullen 1994). Large-scale purse seine operations for Jack Mackerel continued through the 1990s. However, purse seine operations ceased in 2000 due to large inter-annual fluctuations in catches and an overall downward trend in fishery production.

Mid-water trawling to target sub-surface schools of Jack Mackerel off Tasmania was trialled in 2001/02. Between December 2001 and April 2002, a total catch of over 5,000 t of small pelagic fishes was taken, with 90% being Redbait. A multi-purpose 50 m mid-water trawler was used to target small pelagic fishes from late 2002 onwards. By mid-2003, more than 7,000 t of small pelagic fishes had been taken, with Redbait dominating the catch. Trawl effort declined in the late 2000s, whereas small-scale purse seine operations continued into the early 2010s (Emery et al. 2015).

The long-term patterns of production in the fishery for Jack Mackerel off eastern Australia are likely to be the result of a combination of changes in fish availability/abundance and market/economic factors. Several authors have documented large inter-annual variability in oceanographic conditions in the southern part of the East Australian Current (e.g. Harris et al. 1992, Young et al. 1993, McLeod et al. 2012), which may have contributed to changes in relative abundance of surface schools of small pelagic species such as Jack Mackerel and their availability to the fishery. The apparent shift from Jack Mackerel to Redbait as the dominant small pelagic fish in this region during the 1990s may have resulted from changes in food availability caused by environmentally-driven changes in the plankton assemblage (Harris et al. 1992, Young et al. 1993, McLeod et al. 2012). However, the potential effects of fishing on abundance and population structure are still poorly understood.

#### 2.1.2 Taxonomy

Jack Mackerel belong to the family *Carangidae*, which includes 140 species representing 32 genera (Nelson 2006). Carangids are found worldwide with most species occurring in tropical waters. There are 65 species in Australian waters; eight species from four genera inhabit temperate waters (Gomon et al. 2008). The genus *Trachurus* contains 13 species;

three of these species are found in Australia: *T. declivis, T. murphyi* and *T. novaezelandiae*.

#### 2.1.3 Distribution

Jack Mackerel are widely distributed throughout coastal waters of southern Australia and New Zealand. In Australia, this species occurs along the southern coast from Shark Bay in Western Australia to Wide Bay in Queensland, including the waters around Tasmania (Gomon et al. 2008). Jack Mackerel is found to depths of 500 m but is most abundant over the continental shelf to 200 m (Pullen 1994).

#### 2.1.4 Stock Structure

There is some evidence to suggest that at least two populations of Jack Mackerel occur within Australian waters, whilst a third occurs in New Zealand. Analysis of morphometric measurements and meristic counts showed a significant difference between east Australian fish and those from the Great Australian Bight (GAB) (Lindholm and Maxwell 1988). Genetic studies have found no significant differences between southern NSW and eastern Tasmanian populations (Smolenski et al. 1994), but distinct differences between those from the GAB and New Zealand (Richardson 1982).

Mounting evidence, resulting from spawning biomass surveys, indicate a possible East-West break in stock structure along the Bonney Coast of South Australia (Ward et al. 2015a, 2018, 2020a). The main spawning stock for south-eastern Australia is thought to be located from western Victoria through Bass Strait to eastern Tasmania and southern NSW (Bulman et al. 2015, Ward et al. 2018, 2020a).

#### 2.1.5 Movement

No specific studies have examined the movement of Jack Mackerel. However, a correlation between size and depth is evident, with smaller fish generally found inshore and larger fish offshore (Shuntov 1969, Stevens et al. 1984, Kailola et al. 1993, Pullen 1994). Such size-dependent distribution suggests offshore movement with increasing size.

#### 2.1.6 Food and Feeding

Jack Mackerel feed primarily on aquatic crustaceans (Shuntov 1969, Stevens et al. 1984, Bulman et al. 2008, McLeod et al. 2012), and krill (*Nyctiphanes australis*) are the most common dietary item throughout the fish's distribution. Krill accounts for ~44% of the diet in Jack Mackerel from eastern Tasmania (Webb 1976, Williams and Pullen 1986, McLeod et al. 2012). Jack Mackerel living in deeper waters also feed on mesopelagic fish (Maxwell 1979, Blaber and Bulman 1987). In addition, Jack Mackerel eat minor quantities of other prey items, including ostracods, gastropods, amphipods, isopods, polychaetes and echinoderms (Stevens et al. 1984, Blaber and Bulman 1987, McLeod et al. 2012). Dietary composition also varies seasonally (Bulman et al. 2008). In the GAB, Jack Mackerel generally feed during the day with fish in offshore waters feeding mostly on krill and fish in inshore waters consuming mainly copepods (Shuntov 1969, Stevens et al. 1984). Prey size is dependent on fish size, with larger prey items taken by larger fish (Stevens et al. 1984).

#### 2.1.7 Age, Growth and Size

Jack Mackerel reach a maximum of 470 mm FL, 1 kg in weight and 17 years of age (Last et al. 1983, Williams and Pullen 1986, Lyle et al. 2000, Browne 2005). Multiple studies have investigated the age and growth of Jack Mackerel (whole otoliths: Stevens and Hausfeld 1982, Jordan 1994; sectioned otoliths: Lyle et al. 2000, Browne 2005). The annual formation of increments in otoliths has been validated using bomb radiocarbon analysis (Lyle et al. 2000). In Tasmania, Jack Mackerel grow quickly at a young age, reaching 270 mm total length (TL) within their first 4 years and 335 mm TL by 10 years, with no significant difference in growth between males and females (Lyle et al. 2000). Differences in growth rates have been reported for Jack Mackerel off Kangaroo Island compared to those off southern NSW; fish off Kangaroo Island were larger at a given size than those off southern NSW (Dennis et al. 2021).

### 2.1.8 Reproduction

Jack Mackerel are serial spawners (Marshall et al. 1993, Neira 2011). Fish spawn in spring along the NSW coast (Maxwell 1979, Keane 2009) and during summer further south off Tasmania and in the Great Australian Bight (Stevens et al. 1984, Jordan et al. 1995, Ward et al. 2016, Sexton et al. 2017). The main spawning area is thought to be located off south-eastern Australia from western Victoria through Bass Strait to eastern Tasmania and southern NSW (Bulman et al. 2015, Ward et al. 2018, 2020a). Off eastern Tasmania, spawning occurs continuously from December to February and follows a semi-lunar cycle, where peak activity is associated with both full and new moons (Williams and Pullen 1986, Jordan 1994, Neira 2011). Mean gonadosomatic index (GSI) values for females off eastern Tasmania increase substantially in November and remain high until January, before declining in February (Williams and Pullen 1986; Ward et al. 2011b, 2013).

Fifty percent sexual maturity occurs around 270 mm fork length (FL) for both sexes (Webb 1976, Ward et al. 2011b, 2013). During recent DEPM surveys for Jack Mackerel East, the mean size of mature females collected was 266 mm FL in 2014 (Ward et al. 2015a) and 228 mm FL in 2019 (Ward et al. 2020a). Relative fecundity has been estimated to be ~200 oocytes per gram of weight for fish off southern NSW to south-eastern Tasmania (Ward et al. 2020a). Mean spawning fraction (proportion of mature females spawning per day/night) is estimated to range from 0.032 to 0.056 in southern Australia (Ward et al. 2015a, 2016, 2020a).

#### 2.1.9 Early Life History and Recruitment

Jack Mackerel eggs are positively buoyant and 0.97–1.03 mm in diameter (Neira 2011). Larvae have been described in Neira et al. (1998). Larvae have been collected off southern NSW during spring, and off eastern Tasmania, in Bass Strait and the GAB during summer (Stevens et al. 1984, Keane 2009, Ward et al. 2015a, 2018, 2020a). Jack Mackerel eggs are morphologically similar to Yellowtail Scad eggs but slightly larger (Yellowtail Scad egg diameter: 0.78–0.88 mm; Neira 2009).

#### 2.1.10 Stock Assessment

During the late 1980s and early 1990s, considerable research effort was directed at describing the fisheries biology of Jack Mackerel. Projects were initiated to: (1) evaluate tools for assessment of stocks; (2) describe factors contributing to inter-annual variability in abundance; and (3) collect information on early life history and reproductive biology (Jordan et al. 1992, 1995). Research outputs included greater understanding of interactions between local oceanography and presence of surface schools of Jack Mackerel (Harris et al. 1992, Williams and Pullen 1993), and data on their reproductive biology and early life history (Harris et al. 1992, Marshall et al. 1993, Williams and Pullen 1993, Jordan 1994, Jordan et al. 1995). The abundance of surface schools off eastern Tasmania was closely related to oceanographic changes (Young et al. 1993). However, no successful method of assessing the size of the Jack Mackerel resource was developed, despite attempts to use a combination of aerial surveys of surface-schooling fish and hydro-acoustic surveys of surface and sub-surface schools on the shelf break (Jordan et al. 1992).

Dedicated DEPM surveys for Jack Mackerel East were conducted in 2014 and 2019 (Ward et al. 2015a, 2020a). A preliminary DEPM was done in 2011 using samples collected off south-eastern Australia in 2002–2004 during a survey for Blue Mackerel (Neira 2011). Ecosystem modelling of south-east Australian waters has also been used to estimate the spawning biomass of Jack Mackerel (Fulton 2013). The first dedicated DEPM survey for Jack Mackerel along the southern Australian coast (Kangaroo Island to western Tasmania) occurred during December 2016 and January 2017 (Ward et al. 2018).

#### 2.1.11 Recreational fishing

In Australia, recreational fishers target Jack Mackerel using rod and line, and troll lines in NSW, Queensland, South Australia, Western Australia and Tasmania. The Australian National Survey of Recreational and Indigenous Fishing (Henry and Lyle 2003) estimated that boat-based recreational fishers harvested 740,260 individual Jack Mackerel and Scads (combined) in 2000/01, with 37% of these being released back into the water. Of those fish retained, 46% were taken in NSW, 26% in Western Australia and 19% in Queensland (Henry and Lyle 2003). Based on the mean length/weight key developed by Stewart and Ferrell (2001), the estimated weight of Jack Mackerel/Yellowtail Scad harvested by the recreational sector annually in Australia was ~94 t (Ward and Rogers

2007). This catch information is not presented in this report, as estimates of catch for individual species were not available.

#### 2.1.12 Biomass Estimates

#### East

The DEPM has been applied to Jack Mackerel East in 2002 (Neira 2011), 2014 (Ward et al. 2015a) and 2019 (Ward et al. 2020a). The preliminary application of the DEPM estimated the spawning biomass of Jack Mackerel East between Sugarloaf Point and Cape Howe, NSW during October 2002 to be 114,000–169,000 t (Neira 2011). This estimate was considered imprecise due to a lack of samples to estimate adult reproductive parameters.

The first dedicated application of the DEPM to Jack Mackerel East was undertaken in January 2014 off south-eastern Australia; the estimate of spawning biomass for the survey was 157,805 t (95% CI = 59,570–358,731 t) (Ward et al. 2015a). The estimated spawning area off south-eastern Australia was 23,553 km<sup>2</sup>, comprising 37.2% of the total area sampled (63,355 km<sup>2</sup>) (Ward et al. 2015a). Live Jack Mackerel eggs (n = 3,530) were collected from 117 of the 292 (40.1%) stations between Port Stephens, NSW and South East Cape, Tasmania. Mean daily egg production ( $P_0$ ) was 28.9 eggs·day<sup>-1</sup>·m<sup>-2</sup>. The highest densities of Jack Mackerel eggs were recorded in waters off north-eastern Tasmania and in Bass Strait, and SSTs ranged between 15 and 22°C.

The second dedicated application of the DEPM to Jack Mackerel East was undertaken in January/February 2019 off south-eastern Australia; the estimate of spawning biomass for the survey was 156,292 t (95% CI = 49,120–263,496 t) (Ward et al. 2020a). The estimated spawning area off south-eastern Australia was 36,100 km<sup>2</sup>, comprising 52.9% of the total area sampled (68,295 km<sup>2</sup>) (Ward et al. 2020a). Live Jack Mackerel eggs (n = 921) were collected from 107 of the 206 (51.9%) stations between Jervis Bay, NSW and South East Cape, Tasmania.  $P_0$  was 15.1 eggs·day<sup>-1</sup>·m<sup>-2</sup>. The highest densities of Jack Mackerel eggs were recorded in Bass Strait and off Batemans Bay; SSTs ranged from 18 to 23°C.

Sensitivity analyses showed that realistic variations of each parameter produced estimates of spawning biomass for Jack Mackerel East that were between about 100,000 t and 200,000 t. Variations in the values of spawning fraction and  $P_0$  caused the greatest change in estimates of spawning biomass (Ward et al. 2015a, 2020a).

The spawning biomass of Jack Mackerel East appears to have remained stable between 2014 and 2019, despite recent increases in annual catches. The estimates of spawning biomass for 2014 and 2019 are considered to be reliable, as they are based on robust and/or conservative estimates of all key parameters.

#### West

The first dedicated application of the DEPM to Jack Mackerel West was undertaken in December 2016 and January 2017 off southern Australia (Ward et al. 2018). Two major areas of spawning activity—split by the Bonney Coast—were identified: an area south of Kangaroo Island and an area between King Island and the western Victorian coast. In addition to egg samples collected in the main survey area, opportunistic samples were also taken in Bass Strait where a large amount of spawning activity was detected. The estimate of spawning biomass derived from all samples (main survey + Bass Strait) was 31,069 t (Ward et al. 2018). The total estimated spawning area was 13,898 km<sup>2</sup>, comprising 15.7% of the total area sampled (88,374 km<sup>2</sup>) (Ward et al. 2018). Live Jack Mackerel eggs (n = 639) were collected from 55 of the 347 (15.9%) stations between western Kangaroo Island, South Australia and south-western Tasmania.  $P_0$  was 9.6 eggs·day<sup>-1</sup>·m<sup>-2</sup>. The highest densities of Jack Mackerel eggs were recorded in waters off north-west of King Island (TAS), south-east of Kangaroo Island and in Bass Strait; SSTs ranged between 16 and 20°C.

The estimate of spawning biomass derived from all samples is indicative of adult abundance between western Kangaroo Island and south-western Tasmania, but is an underestimate of the total abundance of Jack Mackerel in the West sub-area. Bass Strait was not sampled extensively, and the western Jack Mackerel stock is known to extend west of Kangaroo Island (e.g. Stevens et al. 1984, Bulman et al. 2015). Sensitivity analysis showed that, for this study, spawning area had a larger effect on estimates of spawning biomass than S or  $P_0$  (Ward et al. 2018).

#### 2.1.13 Management Strategy Evaluation

Smith et al. (2015) concluded the harvest rate of 15% may be too high for Jack Mackerel and suggested a Tier 1 harvest rate of 12% for Jack Mackerel East and West, with the Tier 1 rate being applied for not more than 5 years. Tier 2 harvest rates for Jack Mackerel East and West were recommended to be 50% of Tier 1 rates and not to be applied for more than 10 years. The study indicated that it is not safe to apply Tier 2 harvest rates unchecked for long periods of time (i.e. >10 years; Smith et al. 2015). The Atlantis-SPF biomass estimate for Jack Mackerel East used by Smith et al. (2015) was 137,000 t (typical range: 91,000–208,000 t) and 62,000 t (typical range: 60,000–110,000 t) for Jack Mackerel West (Smith et al. 2015).

#### 2.1.14 Management

Currently, Jack Mackerel East and West are managed at the Tier 1 level under the Harvest Strategy. DEPM assessments of Jack Mackerel have been conducted for both the East and West sub-areas of the SPF: Jack Mackerel East in 2002, 2014 and 2019, and Jack Mackerel West in 2016/17.

## 2.2 Methods

#### 2.2.1 Fishery Statistics

Fishery statistics from 1984/85 to 2020/21 were supplied by relevant jurisdictions and collated by the South Australian Research and Development Institute (SARDI) Aquatic Sciences. Unless indicated, annual data are reported in fishing seasons (May 1 to April 30) rather than financial years.

Estimates of total annual catch for Jack Mackerel East include data from the NSW Ocean Fisheries (Hauling, Trap and Line, Trawl), NSW Estuary General Fishery, Victorian Ocean Purse Seine Fishery, Tasmanian Scalefish Fishery and the Commonwealth SPF. For Jack Mackerel West, total annual catch estimates include data from the Tasmanian Scalefish Fishery, Victorian Ocean Purse Seine Fishery, South Australian Marine Scalefish Fishery and Commonwealth SPF. Due to data confidentiality (<6 licence holders annually reporting catch since 2015/16), fishery data from Victoria were not provided and so are not included in total annual catch statistics since 2015/16.

Mean annual catch per unit effort (CPUE) of Jack Mackerel East and West in the Commonwealth SPF is calculated for the gear types of mid-water trawl (tonnes·trawl-hour<sup>1</sup>  $\pm$ SE) and purse-seine (tonnes·net-set<sup>-1</sup>  $\pm$ SE) from 2000/01 to 2020/21. For the 2020/21 fishing season, shots by mid-water trawl of <0.2 hrs were removed from the analyses, as these were deemed 'false shots' (pers. comm. AFMA). Zero catch of Jack Mackerel in a trawl was assumed when fishing occurred but catch of Jack Mackerel was not reported in the logbook record.

#### 2.2.2 Biological Information

Fishery-dependent length frequency and biological data were collected between 1984 and 1993 as part of a monitoring program of the Jack Mackerel Purse Seine Fishery off Tasmania. Samples collected between 1985 and 1990 during demersal research trawling, conducted by CSIRO and the Tasmanian fisheries agency, supplied some biological information. Between 1994 and 2001, the level of catch sampling of the purse seine fishery was limited.

Biological data were collected by AFMA observers on a small proportion of trips during the 2001/02 pair-trawl fishing trials undertaken off Tasmania. When mid-water trawl operations started in 2002, the Tasmanian Aquaculture and Fisheries Institute (TAFI) began an intensive biological monitoring program that continued to 2006. AFMA also provided observer coverage of mid-water trawl operations, with additional length-frequency data collected from 2002 to 2008.

Purse seine operations for small pelagic fish resumed in Tasmanian State waters in 2008/09, mainly targeting Redbait and Jack Mackerel. Catch sampling was implemented in 2009/10 as part of the SPF monitoring program under the Harvest Strategy (AFMA 2008).

No catch samples were obtained for Jack Mackerel from 2010/11 to 2013/14 due to limited fishing activity. Catch sampling by AFMA observers resumed in the SPF in 2014/15. Samples of Jack Mackerel were collected (n = 50 randomly selected fish per trawl) and supplied to SARDI Aquatic Sciences to estimate the current size and age composition of the catch.

Biological data collected from each fish include: body length (mm FL), total weight (g), sex, gonad developmental stage (following the macroscopic staging criteria described in Marshall et al. 1993) and gonad weight (to the nearest 0.1 g). Gonad stages were designated as: I) immature; II) maturing virgins or recovering spent; III) maturing; IV) ripe; and V) spent. Otoliths were removed from random sub-samples of fish for age estimation. The age structure of Jack Mackerel prior to 2014/15 was estimated using age-length keys based on age data pooled from 1985/86, 1989/90, 1993/94 and 1994/95. Since 2014/15, ages for Jack Mackerel have been based on annual growth increment counts in thinsectioned otoliths (sub-samples of 5 to 10 fish per sample).

Jack mackerel otoliths were aged following the protocol of Lyle et al. (2000) after thinsectioning and mounting on microscope slides. Edges of each annual opaque zone were counted because they are more easily delineated than edges of translucent zones. The edge of the first annual opaque zone was identified using guidelines described of Lyle et al. (2000), where a sub-annual inner increment was sometimes present and was disregarded. The period of age represented by the first annual increment was assumed to be similar across all regions where the fish were collected, based on spawning stock, spawning time and time of opaque increment formation (Lyle et al. 2000, Ward et al. 2015a, Ward et al. 2016). All fish were assigned a birthdate of 1 January based on the known spawning off southern Australia (Ward et al. 2016). Jack mackerel deposit the opaque regions of their otoliths during winter (Lyle et al. 2000). The completion of annual opaque zones in otoliths during spring/summer in temperate marine waters is common (e.g. Choat and Axe 1996, Fowler and Short 1998, Smith and Deguara 2003, Ewing et al. 2007). A standardised completion date of 31 October was designated for the opague zone for all regions. The first year of growth of Jack mackerel in southern Australia represents growth from 1 January to 31 October (10 months). Each otolith was aged by two different readers and the counts compared. Where the counts differed, the otolith was re-read and an age consensus reached, or the otolith was removed from further analyses. Prior to ageing otoliths, the readers used a species-specific reference collection to calibrate their ageing technique (see Ward and Grammer 2019).

Catch weighting was applied to length/age-frequency data collected since 2014/15. Length- and age-frequencies were weighted by the number of fish sampled per trawl to account for uneven sample sizes and then catch weighted by the total amount of Jack Mackerel taken in the trawl.

Commercial logbook information, length-frequency and biological data collected between 1984 and 2021 are included in this assessment. In addition to current catch samples, age, growth and reproductive data from previous studies are included (Jordan et al. 1992, Lyle

et al. 2000, Browne 2005, Ward et al. 2011b). Length-frequency data from research sampling undertaken in January 2014 (Ward et al. 2015a) and January/Februay 2019 (Ward et al. 2020a) are also included. Summarised biological data prior to 2014/15 are presented in financial years. From 2014/15 to present, all SPF catch sampling data are presented in fishing seasons from 1 May to 30 April.

### 2.3 Results

#### 2.3.1 Jack Mackerel East

### 2.3.1.1 Fishery Statistics

#### Number of vessels

The total number of vessels reporting catches of Jack Mackerel East from all jurisdictions declined from >100 vessels prior to 1998/99 to 11 vessels in 2012/13 and increased to 18 vessels in 2013/14. On average, 14 vessel per year have landed Jack Mackerel in the East sub-area since then. A total of 17 vessels reported catches of Jack Mackerel East in 2018/19, 16 in 2019/20 and 13 in 2020/21. The total number of Commonwealth vessels reporting catches of Jack Mackerel East increased from 1 in 2017/18 to 3 in 2020/21.

#### Annual patterns: Total catch

Total catches of Jack Mackerel East declined from ~40,000 t in 1986/87 to 310 t in 1995/96 (Figure 2-1a). Catches increased to 9,916 t in 1997/98 and decreased through the 2000s with catches of <100 t from 2011/12 to 2013/14. Total catch increased to 6,321 t in 2015/16, dropped to 2,751 t in 2017/18, and rose to 7,912 t in 2019/20 (Figure 2-1a). Total catch was 5,894 t in 2020/21. The main fishery for Jack Mackerel East is the SPF (purse seine and mid-water trawl).

#### Annual patterns: Catch, Effort and CPUE

Within the SPF, purse seining has historically been used to target Jack Mackerel East (Figure 2-1b). Purse seine effort for Jack Mackerel East declined from 15 net-sets in 2004/05 to zero net-sets from 2012/13 to 2019/20 and increased to 2 net-sets in 2020/21 (Figure 2-1b). Annual purse seine catch follows effort with 307 t taken in 2000/01, no catch from 2012/13 to 2019/20 and <2 t in 2020/21 (Figure 2-1b). Mean annual CPUE for purse seining in the SPF has declined from 44 t·net-set<sup>-1</sup> in 2000/01 to <1 t·net-set<sup>-1</sup> in 2020/21 (Figure 2-1c).

Mid-water trawling replaced purse seining in the SPF for Jack Mackerel East after the early 2000s (Figure 2-1d). Annual trends of mid-water trawl effort and catch are similar since 2000/01: both increased in 2003/04 (1,338 trawl-hours; 3,300 t), decreased to zero effort and catch from 2011/12 to 2013/14 and increased to 1,307 trawl-hours in 2019/20 with 7,910 t of catch. Effort increased in 2020/21 to 1,483 trawl-hours while catch declined to 5,888 t (Figure 2-1d). Mean annual CPUE of mid-water trawls increased from 0 t-trawl-

hour<sup>-1</sup> in 2011/12 to 11 t trawl-hour<sup>-1</sup> in 2016/17 and declined to 4 t trawl-hour<sup>-1</sup> in 2020/21 (Figure 2-1e).



Figure 2-1. Fishery statistics for Jack Mackerel East. (A) Total annual landed catch (tonnes) for all jurisdictions from 1984/85 to 2020/21. Black bars: catch per financial year; grey bars: catch per fishing season. Long-term trends in the SPF by fishing season from 2000/01 to 2020/21: (B) annual landed catch (tonnes) and effort (net-sets) by purse seine; (C) mean annual CPUE (t·net-set<sup>-1</sup>;  $\pm$ SE) by purse seine; (D) annual landed catch (tonnes) and effort (trawl-hours) by mid-water trawl; (E) mean annual CPUE (t·trawl-hour<sup>-1</sup>;  $\pm$ SE) by mid-water trawl.

## 2.3.1.2 Biological Information

Length-frequency data were collected from 7,458 Jack Mackerel sampled from commercial mid-water trawl catches off eastern Tasmania between 2001/02 and 2009/10 (Table 2-1). Age-frequency data were obtained from 7,258 of those fish (Table 2-1). Catch samples were not collected from 2010/11 to 2013/14. Catch sampling in the SPF resumed in 2014/15. Between 2014/15 and 2020/21, 243 samples were collected in the SPF from midwater trawl catches in the East sub-area (Table 2-1). Length-frequency data were collected from 9,936 Jack Mackerel. Age-frequency data were obtained from 2,898 of those fish (Table 2-1).

In January 2014, 10 samples (n = 1,759) of Jack Mackerel were collected during DEPM research surveys (demersal trawl net) off eastern Victoria and southern NSW (Ward et al. 2015a). An additional 7 samples (n = 947) were collected from waters off north-eastern Tasmania (Ward et al. 2015a). During the 2019 DEPM survey for Jack Mackerel East (January-February 2019), 12 samples (n = 1,169) of Jack Mackerel were collected in the same region (Ward et al. 2020a).

Season	SPF sub- area	Gear type	No. of samples	Length- frequency (n)	Age- frequency (n)	Size range (mm FL)	Age range (years)
2001/02	East	mid-water trawl	-	-	31	-	2-16
2002/03	East	mid-water trawl	-	687	687	120-380	2-16
2003/04	East	mid-water trawl	-	4,736	4,736	160-390	2-16
2004/05	East	mid-water trawl	-	1,717	1,717	180-350	2-15
2009/10	East	mid-water trawl	5	318	87	150-290	2–7
2014/15	East	mid-water trawl	7	325	235	185-380	2–15
2015/16	East	mid-water trawl	94	4,532	1,197	82-425	1–11
2016/17	East	mid-water trawl	30	1,478	350	144-306	2–10
2017/18	East	mid-water trawl	29	1,438	346	178-291	1–10
2018/19	East	mid-water trawl	22	1,184	240	116-282	0-10
2019/20	East	mid-water trawl	33	618	385	175-255	1-11
2020/21	East	mid-water trawl	28	361	145	149-255	2-10

Table 2-1. Summary of Jack Ma	ackerel East catch samples co	ollected from commercial SPF landings
-------------------------------	-------------------------------	---------------------------------------

#### Size structure

#### The purse seine fishery: 1984/85–2009/10

Purse seine catches of Jack Mackerel East off eastern Tasmania between 1984/85 and 1995/96 mainly contained fish between 210 and 350 mm FL, and included individuals up to 440 mm FL (Figure 2-2). The size structures over those years contained both unimodal (mode range: 250-310) and bimodal distributions (mode range: 230-330). From 1993/94–



1995/96, size structures were unimodal with a shift to larger fish. In 2009/10, the size structure was unimodal and dominated by smaller fish (mode: 200 mm FL; Figure 2-2).

Figure 2-2. Length-frequency distributions of Jack Mackerel East caught in the SPF by purse seine from 1984/85 to 1995/96 and in 2009/10; n = number of fish.

#### Mid-water trawl fishery: 2002/03–2020/21

Jack Mackerel caught by mid-water trawl off eastern Tasmania from 2002/03 to 2004/05 were mostly between 200 and 300 mm FL (Figure 2-3), and considerably smaller than those caught in earlier purse seine operations. The size composition of mid-water trawl and purse seine catches in the East during 2009/10 were similar, with each dominated by fish between 180 and 240 mm FL (Figures 2-2 and 2-3).

Modal length of Jack Mackerel increased from 240 to 270 mm FL between 2002/03 and 2004/05 in mid-water trawl catches taken in the East, but only a small proportion of the catch contained fish >300 mm FL (Figure 2-3). In 2014/15, the size structure of catch samples from mid-water trawling off southern NSW increased compared to 2009/10. The modal length increased to 220 mm FL, and more fish were in the 250 to 300 mm FL size classes (Figure 2-3). The modal length of the 2014/15 commercial mid-water trawl catch was smaller (220 mm FL) than the commercial trawl catches of the early 2000s (240–270 mm FL; Figure 2-3). The modal length increased to 230 mm FL in 2015/16 and 2016/17, then decreased to 200 mm FL in 2017/18 and 2018/19. Modal length increased to 210 mm in 2019/20 and decreased to 190 mm in 2020/21 (Figure 2-3).



Figure 2-3. Length-frequency distributions of Jack Mackerel East caught by mid-water trawl in the SPF from 2002/03 to 2020/21; n = number of fish.

#### Research (demersal trawl net) surveys: 2014 and 2019

The research samples of Jack Mackerel collected off eastern Victoria/southern NSW (above 39°S) in 2014 mainly consisted of fish between 240–310 mm FL, with a mode at 250 mm FL (Figure 2-4). Similarly, the size structure for research samples off north-eastern Tasmania (below 39°S) in 2014 mainly contained fish between 250–310 mm FL, although with a narrower range and stronger mode (280 mm FL; Figure 2-4). In 2019, the size structure for research samples collected from both eastern Victoria/southern NSW (190–230 mm FL; mode: 220 mm FL) and north-eastern Tasmania (210–240 mm FL; mode: 220 mm FL) was smaller than in 2014 (Figure 2-4).

The modal length of Jack Mackerel in research catches from eastern Victoria/southern NSW in 2014 was larger compared to that from commercial trawl catches taken in the same area in 2014/15 (Figures 2-3 and 2-4). In 2019, the modal length from research catches was only slightly larger than that from mid-water trawl catches in 2019/20 (Figures 2-3 and 2-4).





#### Age structure

#### The purse seine fishery: 1984/85–2009/10

Jack Mackerel taken by purse seine off eastern Tasmania were generally aged 3 to 10 years (Figure 2-5). Catches between 1984/85 and 1990/91 were dominated by fish aged 4 to 5 years with fish up to 9 years also well represented. Between 1991/92 and 1994/95, few fish older than 6 years were taken; fish were mainly aged 3 to 5 years. The 1995/96

age structure was similar to that of the mid 1980s suggesting the relative scarcity of older fish in the intervening years may not have been solely due to the impact of fishing on the population age structure. However, it should be noted that using a pooled age-length key rather than annual age data may have had a smoothing effect on age composition, particularly when representing the older age groups. In 2009/10, purse seine catches from eastern Tasmania were mainly fish aged 2 to 3 years (Figure 2-5).



Figure 2-5. Age-frequency distributions of Jack Mackerel East caught in the SPF by purse seine from 1984/85 to 1995/96 and in 2009/10; n = number of fish in distribution.

#### Mid-water trawl fishery: 2001/02–2020/21

Mid-water trawl catches of Jack Mackerel off eastern Tasmania between 2001/02 and 2004/05 were mainly fish aged 2–5 years old, with a modal age of 4 years (Figure 2-6). During 2009/10, mid-water trawl catches from eastern Tasmania mostly contained fish

#### Securing Australia's fishing future

aged 2–3 years. In mid-water trawl catches off southern NSW during 2014/15, 72% of the fish were between 3 and 6 years (age mode: 4 years) (Figure 2-6). In 2015/16, the modal age was 5 years with 70% of fish between 4 and 6 years. In 2016/17, 5–6 year olds made up 55% of the age structure. In 2017/18, the modal age was 4 years with 60% of the fish between 3 and 5 years (Figure 2-6). In 2018/19, 2019/20 and 2020/21 modal age was 3 years with 90% of the fish ≤4 years in 2020/21. The maximum estimated age of Jack Mackerel East from mid-water trawl catch samples was 16 years (Figure 2-6).



Figure 2-6. Age-frequency distributions of Jack Mackerel East caught by mid-water trawl in the SPF from 2001/02 to 2020/21; n = number of fish in distribution.

#### 2.3.2 Jack Mackerel West

### 2.3.2.1 Fishery Statistics

#### Number of vessels

The total number of vessels reporting catches of Jack Mackerel West from all jurisdictions declined from 17 in 1998/99 to 1 in 2012/13. Total vessel numbers increased to 6 in 2014/15; 5 vessels were fishing in 2017/18 and 3 in 2018/19. Vessel numbers increased again to 13 in 2019/20, with 11 vessels reporting catches in 2020/21 (0 Commonwealth vessels in 2020/21).

#### Annual patterns: Total catch

Historically, Jack Mackerel catches in the West sub-area have been lower than in the East, with the SPF taking 93% of the total annual catch since 1995/96. Due to data confidentiality (<6 licence holders reporting catch per year) only SPF catches are discussed in this section (Figure 2-11a). Annual catches of Jack Mackerel West (purse seine and mid-water trawl) did not exceed 360 t prior to 2015/16 (Figure 2-11a). Catches in the SPF peaked at 359 t in 2006/07 and declined from there onwards, with no catches reported from 2010/11 to 2014/15. Catches increased to 686 t in 2016/17. In 2017/18 and 2018/19, 0 t of Jack Mackerel West was taken in the SPF (Figure 2-11a). In 2019/20 12 t of Jack Mackerel West were reported in the SPF, decreasing to 0 t in 2020/21.

#### Annual patterns: Catch, Effort and CPUE

Mid-water trawls have historically been the main gear type used by the SPF for Jack Mackerel West; purse seines have had limited use (Figure 2-11a). Since 2000/01, annual purse seine effort has not exceeded 7 net-sets (2006/07), and the maximum annual catch was 142 t in 2006/07 (Figure 2-11b). There has been no reported purse seine effort and catch of Jack Mackerel West in the SPF from 2010/11 to 2019/20; 2 net-sets (catch: 2 t) were reported in 2020/21 (Figure 2-11b). Mean annual CPUE of purse seining is similar to effort and catch: peaks in 2006/07 (20 t·net-set<sup>-1</sup>) and 2009/10 (19 t·net-set<sup>-1</sup>), with minimal fishing in other years (Figure 2-11c).

Mid-water trawl effort and catch of Jack Mackerel West peaked at 625 trawl-hours and 232 t during the mid-2000s and decreased to zero effort and catch from 2010/11 to 2013/14 (Figure 2-11d). Fishing effort resumed in 2014/15 (19 trawl-hours; zero catch) and 686 t were taken during 2016/17 in 365 trawl-hours (Figure 2-11d). There has been limited effort and catch of Jack Mackerel West since then and no effort and catch in 2020/2021.

Mean annual CPUE of mid-water trawls in the SPF decreased from 1 t·trawl-hour<sup>-1</sup> in 2001/02 to <1 t·trawl-hour<sup>-1</sup> in 2006/07, peaked in 2009/10 (2 t·trawl-hour<sup>-1</sup>) with no further activity until 2015/16 (1 t·trawl-hour<sup>-1</sup>). CPUE increased to 2 t·trawl-hour<sup>-1</sup> in 2016/17, decreased to 0 t·trawl-hour<sup>-1</sup> in 2017/18 and was <1 t·trawl-hour<sup>-1</sup> in 2019/20 (Figure 2-11e).



Figure 2-11. Fishery statistics for Jack Mackerel West. (A) Total annual landed catch (tonnes) for all jurisdictions from 1995/96 to 2020/21. (\*) indicates data confidentiality and only catches by the SPF are shown. Long-term trends in the SPF by fishing season from 2000/01 to 2020/21: (B) annual landed catch (tonnes) and effort (net-sets) by purse seine; (C) mean annual CPUE (t-net-set<sup>-1</sup>; ±SE) by purse seine; (D) annual landed catch (tonnes) and effort (trawl-hours) by mid-water trawl; (E) mean annual CPUE (t-trawl-hour<sup>-1</sup>; ±SE) by mid-water trawl.

## 2.3.2.2 Biological Information

Length-frequency data were collected from 407 Jack Mackerel sampled from commercial mid-water trawl catches off south-western Tasmania between 2002/03 and 2009/10 (Table 2-4). Age-frequency data were obtained from 295 of those fish (Table 2-1). Catch samples

were not collected from 2010/11 to 2014/15 and in 2020/21 due to low levels of fishing activity in the West sub-area. Between 2015/16 and 2019/20, 39 samples of Jack Mackerel were collected in the SPF from midwater trawl catches in the West sub-area (Table 2-4). Length-frequency data were collected from 1,887 Jack Mackerel. Age-frequency data were obtained from 579 of those fish (Table 2-4).

 Table 2-2. Summary of Jack Mackerel West catch samples collected from commercial SPF landings since

 2009.

Season	SPF sub- area	Gear type	No. of samples	Length- frequency n	Age- frequency n	Size range (mm FL)	Age range (years)
2002/03	West	mid-water trawl	-	109	109	260-380	2-16
2003/04	West	mid-water trawl	-	102	102	250-370	2-16
2004/05	West	mid-water trawl	-	64	64	260-350	2-15
2009/10	West	mid-water trawl	1	132	20	160–220	1–3
2015/16	West	mid-water trawl	14	670	281	211–355	2–10
2016/17	West	mid-water trawl	24	1,197	286	228–395	4–13
2019/20	West	mid-water trawl	1	20	12	218-254	3-5

#### Size structure

#### The purse seine fishery

Size structure data are not available for purse seine catches of Jack Mackerel West.

#### Mid-water trawl fishery: 2002/03–2019/20

Jack Mackerel caught by mid-water trawl operations off south-western Tasmania from 2002/03 to 2004/05 were mainly between 250 and 370 mm FL, with an overall modal length of 290 mm FL (Figure 2-12, Table 2-3). Jack Mackerel taken in the West sub-area over this period were larger than those from the East (East overall modal length: 260 mm FL) (Figures 2-3 and 2-12).

In 2009/10, a sample from a single mid-water trawl catch from the West (south-western Tasmania) contained Jack Mackerel of similar sizes (modal length: 190 mm FL) to catches from eastern Tasmania (modal length: 210 mm FL) (Tables 2-1and 2-3; Figure 2-3 and 3-12). The modal length ranged between 240–250 mm FL in 2015/16 and increased to 270 mm FL in 2016/17. Modal length decreased to 230 mm FL in 2019/ 20, but the sample size was small (Figure 2-12).



Figure 2-12. Length-frequency distributions of Jack Mackerel West caught by mid-water trawl in the SPF from 2002/03 to 2019/20; n = number of fish in distribution.

#### Age structure

#### The purse seine fishery

Age structure data are not available for purse seine catches of Jack Mackerel West.

#### Mid-water trawl fishery: 2002/03–2019/20

Annual age structures of Jack Mackerel West sampled from catches taken off southwestern Tasmania from 2002/03 to 2004/05 had a high proportion of fish >5 years, and a mode of 4–5 years (Figure 2-13). The age structure of the single catch in 2009/10 (Table 2-3) contained a high proportion of 2 year old fish, representing about 70% of the distribution (similar to the 2009/10 age structure in the East; Figure 2-3). In 2015/16, the modal age was 4 years with 75% of fish between 3 and 5 years (Figure 2-13). The modal age increased to 7 years in 2016/17, and 70% of fish were 7–8 years of age. Modal age decreased to 3 years in 2019/20 and older fish sampled was 5 years, but the sample size was small (Figure 2-13).

The maximum estimated age of Jack Mackerel West from mid-water trawl catches was 16 years (Figure 2-13, Table 2-3).



Figure 2-13. Age-frequency distributions of Jack Mackerel West caught by mid-water trawl in the SPF from 2002/03 to 2019/20; n = number of fish in distribution.

## 2.4 Summary and Conclusions

#### 2.4.1 Jack Mackerel East

The main fishery for Jack Mackerel East is the SPF; recent catches in other fisheries in this sub-area have been negligible (<15 t per annum). The total annual catch of Jack Mackerel East from 2010/11 to 2014/15 was <250 t. The factory trawler operating in offshore waters from 2014/15 to 2016/17 was replaced in 2017/18 by a mid-water trawler operating in inshore waters off southern NSW. The SPF catch since 2018/19 has been similar to or higher than the catch landed by the factory trawler in 2015/16 and 2016/17. CPUE of Jack Mackerel East since 2017/18 has averaged about 5 t·trawl-hour<sup>-1</sup>, about half that of the factory trawler (10 t·trawl-hour<sup>-1</sup> in 2015/16 and 11 t·trawl-hour<sup>-1</sup> in 2016/17). The reduced CPUE of Jack Mackerel East since 2017/18 compared to 2015/16 and 2016/17 may reflect differences in vessels and gear. Catches reached an all-time high in 2019/20 at 7,910 t from 1,307 trawl-hour<sup>-1</sup> (CPUE of 6 t·trawl-hour<sup>-1</sup>), effort increased to 1,483 trawl-hour<sup>-1</sup> in 2020/21, but catches decreased to 5,888 t, reducing CPUE to <4 t·trawl-hour<sup>-1</sup>.

The modal length of Jack Mackerel taken in 2020/21 was 190 mm FL, below the size at 50% maturity of ~230 mm FL, and down from 210 mm FL in 2019/20 and 200 mm FL in 2018/19. In 2020/21 the modal age was 3 years, similar to modal ages in 2019/20 and 2018/19, but lower than the modal age of 4 years in 2017/18 and 2016/17. The smaller size and younger age of Jack Mackerel taken from East sub-area in 2020/21 may reflect changes in the vessels and gear operating in the fishery.

The average annual total catch of Jack Mackerel East over the last three fishing seasons was 6,251 t. The spawning biomass of Jack Mackerel East in 2019 was estimated to be 156,292 t (95% CI = 49,120–263,496 t) (Ward et al. 2020a). The recent annual average catch of Jack Mackerel East has been 4% of the estimate of spawning biomass and below the Tier 1 exploitation rate for this stock of 12% (Smith et al. 2015). The total catch in 2020/21 in the SPF was 32% of the available TAC. On the basis of the information provided here, Jack Mackerel East is classified as sustainable. Piddocke et al. (2021) also found the biological stock of Jack Mackerel East to be sustainable. Patterson et al. (2021) classified Jack Mackerel East as 'not overfished' and 'not subject to overfishing'.

#### 2.4.2 Jack Mackerel West

The main fishery for Jack Mackerel West is the SPF. Annual catches did not exceed 370 t prior to 2015/16. There was no catch of Jack Mackerel West by the SPF in 2017/18 and 2018/19; previous catches were 686 t in 2016/17 and 634 t in 2015/16. Catches increased in 2019/20 to 12 t, with no catch in 2020/21. The modal length in 2016/17 was 270 mm FL, larger than the mean size at 50% maturity of ~230 mm FL, and larger than the modal length of 240–250 mm FL in 2015/16. Modal length decreased to 230 mm FL in 2019/20, but sample sizes were small. The modal age in 2016/17 was 7 years, up from 4 years in 2015/16, decreasing to 3 years in 2019/20, although sample size was small.

Recent annual catches of Jack Mackerel West have not exceeded 700 t. The spawning biomass of Jack Mackerel West in 2016/17 was estimated to be at least 31,000 t (Ward et al. 2018). Recent catches of Jack Mackerel Wes have been <3% of this estimate of spawning biomass and below the Tier 1 exploitation rate for this stock of 12% (Smith et al. 2015). The total catch in 2020/21 was <1% of the available RBC. On the basis of the information provided here, Jack Mackerel West is assessed as sustainable. Piddocke et al. (2021) also found the biological stock of Jack Mackerel West to be sustainable. Patterson et al. (2021) classified Jack Mackerel West as 'not overfished' and 'not subject to overfishing'.

# **3 Blue Mackerel (Scomber australasicus)**

## 3.1 Introduction

#### 3.1.1 Background to Fishery

Large fisheries for *Scomber* spp. (i.e. ~50,000 to 500,000 t per annum) are located off Japan, Peru, China, Korea, Russia, and the Ukraine (Ward et al. 2001a). The largest fishery for Blue Mackerel in the southern hemisphere is based in New Zealand where annual catches have ranged between ~6,000 t and 15,000 t per annum since the early 1990s (Fisheries New Zealand 2021, MPI 2022). In Australia, Blue Mackerel is taken in several fisheries, and prior to 2018, total annual catches were typically <3,500 t (Ward and Grammer 2019, 2021a). Total annual catches have averaged about 5,000 t since 2018 (Ward and Grammer 2021a).

The NSW commercial purse seine fishery has targeted Yellowtail Scad and Blue Mackerel since the early 1980s (Stewart and Ferrell 2001). During that time, Blue Mackerel typically comprised ~38% of the total annual catches. The average annual catch of Blue Mackerel in Victorian waters between 1978/79 and 2004/05 was 49 t (±22.9 t) with catches ranging between 0.2 and 370.6 t per annum (Ward and Rogers 2007). Blue Mackerel are also an important target species for recreational fisheries in Australia (Henry and Lyle 2003, Giri and Hall 2015, Murphy et al. 2020).

The Tasmanian Purse Seine Fishery has recorded catch and effort data since its inception in 1984. Logbooks contain a record of fishing operations and species taken for each netset. Landings of Blue Mackerel were first reported in 1985/86 with a catch of 587 t (1984/85: 0 t; Pullen 1994). From 1984/85 to 1989/90, Blue Mackerel represented <4% of the total annual catch of small pelagic fishes in Tasmania (Pullen 1994). Species-specific information was not available for other years.

#### 3.1.2 Taxonomy

The genus *Scomber* (family *Scombridae*) historically included three Mackerel species: Blue Mackerel (*S. australasicus*), Chub Mackerel (*Scomber japonicus*), and Atlantic Mackerel (*Scomber scombrus*). However, *S. australasicus* and *S. japonicus* have proved to be more closely related to each other than to *S. scombrus*, and morphological and genetic differences in Atlantic and Indo-Pacific populations of *S. japonicus* warranted recognition of two separate species (Scoles et al. 1998). Atlantic Chub Mackerel (*Scomber colias*) was identified through further genetic analyses and replaces *S. japonicus* in the Atlantic Ocean (Infante et al. 2006, Catanese et al. 2010). Thus, two closely related species occur in the Indian and Pacific Oceans: *S. japonicus* and *S. australasicus*, and two closely related species are found in the Atlantic Ocean: *S. scombrus* and *S. colias*.

### 3.1.3 Distribution

Blue Mackerel occur throughout the Pacific Ocean, including South East Asia, Australia and New Zealand, and in coastal and continental shelf waters of the northern Indian Ocean and Red Sea (depths up to 200m). In Australia, Blue Mackerel are found in subtropical and temperate waters from Queensland to Western Australia and are the only member of the genus present (Ward et al. 2001a, Gomon et al. 2008). Juveniles and small adults usually live in inshore waters, while larger adults form schools in depths of 40–200 m across the continental shelf (Kailola et al. 1993).

#### 3.1.4 Stock Structure

The stock structure of Blue Mackerel in Australasian waters is uncertain. Significant differences in the morphology of monogenean parasites distinguished fish from Australia and New Zealand (Rohde 1987). However, genetic differences have not been found between Blue Mackerel from Australia and New Zealand using mtDNA RFLP analysis and cytochrome *b* sequencing (Scoles et al. 1998). The Australian east coast and west coast Blue Mackerel populations are thought to be genetically separate stocks (Ward and Rogers 2007, Schmarr et al. 2011). An additional stock in southern Australia has tentatively been identified through differentiation with otolith microchemistry and parasite analyses (Ward and Rogers 2007, Schmarr et al. 2017).

Currently, Blue Mackerel off southern Australia are considered to be made up of two biological stocks (Bulman et al. 2008). These stock designations stem from a review and data synthesis to establish management zones in the SPF (Bulman et al. 2008). The Western stock extends from western Tasmania to southern Western Australia, and the Eastern stock occurs to the east of Bass Strait (AFMA 2008, 2009; Bulman et al. 2008).

#### 3.1.5 Movement

No studies have specifically examined the movement of Blue Mackerel in Australasia.

#### 3.1.6 Food and Feeding

Blue Mackerel are pelagic omnivores, feeding mainly on krill, fish and gelatinous nekton (Bulman et al. 2001, Daly 2007, Bulman et al. 2011). Mackerel (*Scomber* spp.) alter their feeding behaviour and ingestion rates depending on prey size and density (Prokopchuk and Sentyabov 2006, Garrido et al. 2007).

#### 3.1.7 Age, Growth and Size

Age estimation in small pelagic fish can be problematic (Gaughan and Mitchell 2000, Arneri et al. 2011), and Blue Mackerel are no exception (Stewart et al. 1999, Ward and Rogers 2007, Marriott and Manning 2011). Although the otoliths of Blue Mackerel have complex inner microstructures, they have been successfully used to estimate annual ages in both Australia (Stewart and Ferrell 2001, Ward and Rogers 2007, Ward and Grammer
2019) and New Zealand (Marriott and Manning 2011). Juveniles of both sexes grow rapidly and reach ~250 mm fork length (FL) after ~2 years of life (Ward and Rogers 2007). Blue Mackerel reach sizes of up to 440 mm FL in the GAB and are estimated to attain ages of ~11 years (Stevens et al. 1984; Ward and Grammer 2019). Growth rates and trajectories of males and females from waters off South Australia are similar (Ward and Rogers 2007). Off eastern Australia, an opaque zone forms in the otoliths of one-year old fish during winter and is complete by early summer (Stewart et al. 1999).

# 3.1.8 Reproduction

Blue Mackerel are serial spawners, and spawn multiple times over a prolonged spawning season with 50% sexual maturity occurring around 237 mm FL for males and 287 mm FL for females (Ward and Rogers 2007, Rogers et al. 2009). Spawning in southern Australia takes place from summer to early autumn and late winter to spring in NSW (Ward and Rogers 2007, Ward et al. 2015b, 2021a). Mean spawning frequencies range from 2 to 11 days in southern Australia. Mean batch fecundity is ~70,000 oocytes per batch and 134 oocytes per gram of weight (Rogers et al. 2009). Fecundity increases exponentially with fish length and weight. Most of the eggs collected off southern and eastern Australia have been obtained from the mid-shelf. High egg and larval densities are recorded in areas with water depths >50 m and with sea surface temperatures (SST) of 18-22°C (Ward and Rogers 2007, Ward et al. 2015b, 2021a). The location of spawning off southern Australia appears to vary substantially among years. Results of an exploratory survey suggest that the western GAB is an important spawning area. However, this region has not yet been sampled intensively (Ward and Rogers 2007).

# 3.1.9 Early Life History and Recruitment

Blue Mackerel eggs are transparent and spherical, measuring 1.05 to 1.35 mm in diameter. The eggs have a smooth chorion, a prominent unsegmented yolk, and a single oil globule 0.22 to 0.38 mm in diameter (Ward and Rogers 2007, Neira and Keane 2008). Blue Mackerel yolk-sac larvae are <3.2 mm total length (TL) at hatching and metamorphose at lengths of ~23.3 mm TL (Neira et al. 1998).

# 3.1.10 Stock Assessment

An extensive study that included both the East and West sub-areas of the SPF investigated the application of a range of egg-based stock assessment methods for Blue Mackerel and concluded that the species was suitable for assessment using the DEPM (Ward and Rogers 2007, Ward et al. 2009). Dedicated DEPM surveys for Blue Mackerel East were conducted in 2014 and 2019 (Ward et al. 2015b, 2021a). Both the annual and daily egg production methods have been used to estimate the spawning biomass of Atlantic Mackerel (*Scomber scombrus*) in the north-eastern Atlantic Ocean (Gonçalves et al. 2009).

# 3.1.11 Recreational fishing

Recreational fishers harvest Blue Mackerel using rod and line, hand line and troll lines (Ward and Rogers 2007) throughout the waters of southern Australia, including southern Queensland. The Australian National Survey of Recreational and Indigenous Fishing estimated that boat-based recreational fishers harvested 720,814 Blue Mackerel annually, with 21% of these being released back into the water (Henry and Lyle 2003). Of the Blue Mackerel retained, 75% were taken in NSW, and 14% and 8% taken in Western Australia and South Australia, respectively (Henry and Lyle 2003). Based on the length/weight key developed by Stewart and Ferrell (2001), the estimated weight of Blue Mackerel harvested annually by the recreational sector in Australia is 228 t (Ward and Rogers 2007).

#### 3.1.12 Biomass Estimates

#### East

The DEPM has been applied to Blue Mackerel East in 2004 (Ward et al. 2009), 2014 (Ward et al. 2015b) and 2019 (Ward et al. 2021a). The preliminary estimate of spawning biomass for Blue Mackerel East in 2004, calculated from the 'best' estimate of each parameter, was 23,009 t (Ward et al. 2009). 'Minimum' and 'maximum' estimates ranged from 7,565 to 116,395 t. The 'best' estimate of spawning biomass was considered to be conservative due to both the approach used to estimate egg production (i.e. McGarvey and Kinloch 2001) and because the survey most likely occurred outside the peak spawning season in that region (Ward and Rogers 2007).

The first dedicated DEPM survey for Blue Mackerel East was undertaken in August/ September 2014 off eastern Australia; the estimate of spawning biomass for the survey was ~83,300 t (95% CI = 35,100–165,000 t) (Ward et al. 2015b). The estimated spawning area off eastern Australia was 17,911 km<sup>2</sup>, comprising 27.3% of the total area sampled (65,528 km<sup>2</sup>) (Ward et al. 2015b). Live Blue Mackerel eggs (n = 2,330) were collected from 70 of the 262 (26.7%) stations between Sandy Cape, Queensland to just south of Newcastle, NSW. Mean daily egg production ( $P_0$ ) was 35.1 eggs·day<sup>-1</sup>·m<sup>-2</sup>. The highest densities of Blue Mackerel eggs were recorded in waters just north of Coffs Harbour and off Port Stephens where SSTs ranged between 18 and 20°C.

The second dedicated DEPM survey for Blue Mackerel East was undertaken in September 2019 off eastern Australia; the estimate of spawning biomass for the survey was 88,265 t (33,320–143,209 t) (Ward et al. 2021a). The estimate of spawning area off eastern Australia was 20,387 km<sup>2</sup>, which was 32.6% of the total area sampled (62,476 km<sup>2</sup>) (Ward et al. 2021a). Live Blue Mackerel eggs (n = 1,829) were collected at 81 of the 251 sites (32%) between southern Queensland and central NSW.  $P_0$  was 36.5 (18.6–59.3) eggs.day<sup>-1</sup>.m<sup>-2</sup>. High egg densities (>10 eggs.m<sup>-2</sup>) were mostly found off the mid-north coast of NSW, in depths of 24–170 m (mean 75 m) and SSTs of 19–20°C.

The estimates of spawning biomass from 2014 (83,300 t) and 2019 (88,265 t) were based on estimates of adult parameters from South Australia and should be treated with caution (Ward et al. 2015b, 2021a). Adult Blue Mackerel collected as part of the 2019 survey were mature, but relatively small and not actively spawning (Ward et al. 2021a). The spawning biomass of Blue Mackerel East appears to have remained stable or increased between 2014 and 2019, despite recent increases in annual catches. Intensive adult sampling to obtain robust estimates of spawning fraction in the East sub-area is needed; this is critical due to the recent increases in annual catches for Blue Mackerel.

Sensitivity analyses showed that realistic variations of each parameter produced estimates of spawning biomass for Blue Mackerel that were between about 50,000 t and 100,000 t. Variations in the values of spawning fraction and  $P_0$  caused the greatest change in estimates of spawning biomass (Ward et al. 2015b, 2021a). Estimates of egg production and spawning area from 2014 and 2019 are likely to be more robust than previously reported.

## West

The preliminary estimate of spawning biomass for Blue Mackerel West in 2005, calculated from the 'best' estimate of each parameter, was 56,228 t (Ward and Rogers 2007). 'Minimum' and 'maximum' estimates ranged from 10,993 t to 293,456 t. The 'best' estimate of spawning biomass was considered to be conservative due to both the approach used to estimate of egg production (i.e. McGarvey and Kinloch 2001) and because there was evidence to suggest that spawning occurred outside the area surveyed in the West (i.e. in the western Great Australian Bight) (Ward and Rogers 2007).

# 3.1.13 Management Strategy Evaluation

Smith et al. (2015) concluded the harvest rate of 15% may be too low for Blue Mackerel and suggested a Tier 1 harvest rate of 23% for Blue Mackerel East and West, with the Tier 1 rate being applied for not more than 5 years. Tier 2 harvest rates for Blue Mackerel East and West were recommended to be 50% of Tier 1 rates and not to be applied for more than 5 years. The study results also indicated it is not safe to apply Tier 2 harvest rates unchecked for long periods of time, particularly on shorter lived species such as Blue Mackerel (Smith et al. 2015). The Atlantis-SPF biomass estimate for Blue Mackerel East is 52,600 t (typical range: 44,000–60,000 t) and 42,500 t (typical range: 34,000–46,000 t) for Blue Mackerel West (Smith et al. 2015).

#### 3.1.14 Management

Currently, Blue Mackerel East is managed at the Tier 1 level, and Blue Mackerel West is managed at the Tier 3 level under the Harvest Strategy. DEPM assessments of Blue Mackerel have been conducted for both the East and West sub-areas of the SPF: Blue Mackerel East in 2004, 2014 and 2019, and Blue Mackerel West in 2005.

# 3.2 Methods

# 3.2.1 Fishery Statistics

Fishery statistics from 1984/85 to 2020/21 have been supplied by relevant jurisdictions and collated by SARDI Aquatic Sciences. Annual data are reported in fishing seasons (May 1 to April 30) rather than financial years as was done in previous assessments (e.g. Ward et al. 2013, 2014c, 2015c).

Estimates of total annual catch for Blue Mackerel East include data from the NSW Ocean Fisheries (Hauling, Trap and Line, Trawl), NSW Estuary Fisheries (General and Prawn Trawl), Tasmanian Scalefish Fishery, Victorian Ocean Purse Seine Fishery and the Commonwealth SPF. In the West, total annual catch estimates include data from the Tasmanian Scalefish Fishery, Victorian Ocean Purse Seine Fishery, South Australian Marine Scalefish Fishery and Commonwealth SPF. Due to data confidentiality (<6 licence holders annually reporting catch since 2015/16), fishery data from Victoria were not provided and have not been included in total annual catch statistics since 2015/16.

Mean annual CPUE of Blue Mackerel East and West in the Commonwealth SPF is calculated for the gear types of mid-water trawl (tonnes·trawl-hour<sup>-1</sup>±SE) and purse-seine (tonnes·net-set<sup>-1</sup>±SE) from 2000/01 to 2020/21. For the 2020/21 fishing season, shots by mid-water trawl of <0.2 hrs were removed from the analyses, as these were deemed 'false shots' (pers. comm. AFMA). Zero catch of Blue Mackerel in a trawl was assumed when effort but not catch was reported in the logbook record.

#### 3.2.2 Biological Information

Length-frequency data for Blue Mackerel East were collected from commercial purse seine catches taken off NSW between 2006/07 and 2014/15. These data were supplied by the NSW Department of Primary Industries (DPI) and are presented in financial years.

Mid-water trawl operations resumed in the SPF during 2014/15; catch sampling required under the Harvest Strategy (AFMA 2008) also recommenced. Samples of Blue Mackerel from mid-water trawl catches (n = 50 randomly selected fish per trawl) were supplied to SARDI Aquatic Sciences to estimate the size and age composition of the catch.

Samples of Blue Mackerel West prior to 2014/15 were obtained from catches taken in summer/early autumn by the commercial purse seine fishery operating from Port Lincoln, South Australia between 2008/09 and 2010/11.

Biological data collected from each fish included body length (mm FL), total weight (±1 g), sex, gonad developmental stage (following the macroscopic staging criteria described in Ward and Rogers 2007) and gonad weight (±0.1 g). Gonad stages were designated as: I) immature; II) maturing virgins or recovering spent; III) maturing; IV) ripe; and V) spent (full descriptions in Ward and Rogers 2007). Otoliths were removed from random sub-samples of fish for age estimation. An otolith weight-age algorithm developed by Ward and Rogers

(2007) was used to estimate ages of Blue Mackerel West prior to 2014/15. Since 2014/15, all Blue Mackerel ages are based on annual growth increment counts in thin-sectioned otoliths (sub-samples of 5 to 10 fish per sample).

Blue mackerel otoliths were aged following the protocol of Marriott and Manning (2011) after thin-sectioning and mounting on microscope slides. Edges of each annual opaque zone were counted (edges of opague zones are more easily delineated than edges of translucent zones). The edge of the opaque core was designated as the first opaque zone. The period of age represented by the opaque core differed based on the region where the fish were collected. All fish collected in southern Australia (SPF sub-zones 3-4) were assigned a birthdate of Jan 1 based on the spawning times for southern Australia (Rogers et al. 2009). In eastern Australia (SPF sub-zones 5 and 6), Blue Mackerel were assigned a birthdate of Sept 1 based on spawning times for eastern Australia (Rogers et al. 2009, Ward et al. 2015b). Blue mackerel form an opaque zone in their otoliths in the winter, but it does not become discernible until spring-summer, i.e. until sufficient translucent material is deposited (Stewart et al. 1999). Therefore, we designated a standardised completion date of Oct 31 for the opaque zone for all regions. The completion of annual opaque zones in otoliths during spring/summer in temperate marine waters is common (e.g. Choat and Axe 1996, Fowler and Short 1998, Smith and Deguara 2003, Ewing et al. 2007). The opaque core (first year of growth) in fish caught in southern Australia represents growth from 1 Jan to 31 Oct (10 months), while the opaque core in fish caught in eastern Australia represents growth from 1 Sept to Oct 31 (14 months). Each otolith was aged by two different readers and the counts compared. Where the counts differed, the otolith was re-read and an age consensus reach, or the otolith was removed from further analyses (see Ward and Grammer 2019).

Catch weighting was applied to length- and age-frequency data collected since 2014/15 in each sub-area. Length- and age-frequencies were weighted by the number of fish sampled per trawl to account for uneven sample sizes and then were catch weighted by the total amount of Blue Mackerel taken in the same trawl.

Biological data prior to 2014/15 are presented in financial years. From 2014/15 to present, all SPF catch sampling data are presented in fishing seasons from 1 May to 30 April.

# 3.3 Results

# 3.3.1 Blue Mackerel East

# 3.3.1.1 Fishery statistics

#### Number of vessels

The total number of vessels reporting catches of Blue Mackerel East from all jurisdictions ranged from 233 to 462 between 1984/85 and 2008/09. Since then, vessel numbers decreased and have ranged between 122 and 162 (149 vessels in 2020/21). On average, 96% of the vessels reporting catch in each year since 2000/01 are from NSW and about 1% are Commonwealth vessels.

#### Annual patterns: Total catch

Total catches of Blue Mackerel East declined from ~1,400 t in 1986/87 to ~100 t in 1989/90 (Figure 3-1). Between 2003/04 and 2011/12, catches ranged from 309 to 1,045 t. From 2012/13 to 2014/15, catches averaged 486 t. Total catch increased to 6,209 t in 2019/20 and reached a historical high of 6,697 t in 2020/21 (Figure 3-1a). The main fisheries that take Blue Mackerel East are the NSW Ocean Hauling Fishery (purse seine) and the SPF (purse seine and mid-water trawl). From 1995/96 to 2014/15, NSW Ocean Fisheries and the SPF took on average 69% and 20% of the total annual catch for the East, respectively. The SPF has taken >80% of the total catch since 2015/16 and >90% since 2019/20.

#### Annual patterns: Catch, Effort and CPUE

Purse seining has historically been used to take Blue Mackerel East. In the SPF, midwater trawling has replaced purse seining (Figure 3-1b, c, e, f), but in the NSW Ocean Hauling Fishery, purse seining is still the primary method used for Blue Mackerel (Figure 3-1d). There has been a long-term decline in purse seine effort in SPF for Blue Mackerel East from a peak of 20 net-sets in 2004/05 to a mean of 2 net-sets annually from 2010/11 to 2015/16 (Figure 3-1b). Effort decreased to zero in 2017/18, increased to 12 net-sets in 2019/20 and 47 in 2020/21. Annual purse seine catch is similar to effort with a high of 280 t in 2005/06,  $\leq$ 1 t taken between 2011/12 and 2017/18, and 17 t in 2020/21 (Figure 3-1b). Mean annual CPUE of purse seines in the SPF decreased from a long term mean of 14 t·net-set<sup>-1</sup> between 2000/01 and 2009/10 to a mean of  $\leq$ 1 t·net-set<sup>-1</sup> from 2010/11 to 2020/21 (Figure 3-1c).



Figure 3-1. Fishery statistics for Blue Mackerel East. (A) Total annual landed catch (tonnes) for all jurisdictions from 1984/85 to 2020/21; black bars: catch per financial year; grey bars: catch per fishing season. Long-term purse seining trends in the SPF by fishing season from 2000/01 to 2020/21: (B) annual landed catch (tonnes) and effort (net-sets); (C) mean annual CPUE (t·net-set-1;  $\pm$ SE). (D) Annual landed catch (tonnes) and effort (net-sets) by purse seining in the NSW Ocean Hauling Fishery. Long-term mid-water trawling trends in the SPF: (E) annual landed catch (tonnes) and effort (ret-sets); (F) mean annual CPUE (t·trawl-hours); (F) mean annual CPUE (t·trawl-hour-1;  $\pm$ SE).

Trends in fishing effort and catch by mid-water trawling in the SPF for Blue Mackerel East are similar (Figure 3-1e). Effort decreased from a peak of 1,338 trawl-hours in 2003/04 (catch: 307 t) to 0 trawl-hours (2011/12–2013/14) and increased to 638 trawl-hours in 2015/16 (catch: 2,022 t). Trawl effort in the SPF fell to 349 trawl-hours in 2016/17 (1,247 t) and has increased to 1,483 trawl-hours in 2020/21 with a catch of 6,202 t (Figure 3-1e). Mean annual CPUE of Blue Mackerel in mid-water trawls in the SPF was <0.4 t·trawl-hour<sup>-1</sup> prior to 2014/15 (Figure 3-1f). CPUE increased to 7 t·trawl-hour<sup>-1</sup> in 2014/15 and has been ~4 t·trawl-hour<sup>-1</sup> since 2018/19 (Figure 3-1f).

# 3.3.1.2 Biological Information

Length-frequency data were collected from 8,887 Blue Mackerel sampled from commercial purse seine catches off NSW between 2006/07 and 2017/18 (Table 3-1). Information on the spatial and temporal coverage of these samples relative to fishery production in NSW was not available. Between 2014/15 and 2020/21, 196 samples were collected in the SPF from midwater trawl catches in the East sub-area (Table 3-2). Length-frequency data were collected from 7,530 Blue Mackerel. Age-frequency data were obtained from 2,363 of those fish (Table 3-2).

Season	SPF sub- area	Gear type	No. of samples	Length- frequency n	Size range (mm FL)
2006/07	East	purse seine	23	1,869	220–400
2007/08	East	purse seine	13	1,286	160–340
2011/12	East	purse seine	13	810	180–390
2012/13	East	purse seine	2	108	280–370
2013/14	East	purse seine	11	1,177	170–360
2014/15	East	purse seine	12	1,382	180–370
2017/18	East	purse seine	31	2,255	210–390

 Table 3-1. Summary of Blue Mackerel samples collected from commercial NSW State catches between

 2006/07 and 2017/18 (data supplied by NSW DPI).

Season	SPF sub- area	Gear type	No. of samples	Length- frequency n	Age- frequency n	Size range (mm FL)	Age range (years)
2014/15	East	mid-water trawl	7	264	221	242–342	2–8
2015/16	East	mid-water trawl	61	2,977	777	187–398	0–11
2016/17	East	mid-water trawl	22	1,090	251	173-351	1–5
2017/18	East	mid-water trawl	26	1,202	405	157-340	0—6
2018/19	East	mid-water trawl	19	1,019	197	146-345	0-9
2019/20	East	mid-water trawl	33	640	389	182-369	1-7
2020/21	East	mid-water trawl	28	338	123	178-314	1-9

Table 3-2. Summary of Blue Macker	I East catch samples collected from	om commercial SPF landings.
-----------------------------------	-------------------------------------	-----------------------------

## Size structure

## Purse seine fishery (NSW): 2006/07–2017/18

There were substantial differences in the size distributions among years of Blue Mackerel East sampled from purse seine catches (Figure 3-2). Fish ranged from 160 to 400 mm FL (Table 3-1). From 2006/07 to 2012/13 (excepting 2007/08), size distributions contained a single mode between 290 and 310 mm FL; the bimodal distribution of 2007/08 had modes at 240 mm FL and 280 mm FL. In 2013/14 and 2014/15, the size distribution included fish from 170 to 370 mm FL, with a single mode at 250 mm FL in each year (Figure 3-2). The length mode increased to 290 mm FL in 2017/18.



Figure 3-2. Length-frequency distributions of Blue Mackerel East collected from purse seine net-sets in NSW between 2006/07 and 2017/18. Data supplied by NSW DPI; n = number of fish. See Table 3-1 for sample N.

#### Mid-water trawl fishery: 2014/15–2020/21

The modal length of Blue Mackerel East from mid-water trawl catch samples from 2014/15 to 2016/17 varied between 260 and 280 mm FL, with fish sizes ranging from 187 to 398 mm FL (Figure 3-3; Table 3-2). There was a bimodal distribution of fish lengths in 2017/18 with a small mode at 170 mm FL and larger one at 270 mm FL; 61% of the fish were 260–280 mm FL (Figure 3-3). The mode increased to 280 mm FL in 2018/19 and dropped to 240 mm FL in 2019/20. In 2020/21, the modal length of fishes was 250 mm FL; 68% measuring between 240 and 260 mm FL. These length-frequency distributions from mid-water trawling are similar to those from purse seine catches taken in the East (Figure 3-2) and 3-3).



Figure 3-3. Length-frequency distribution of Blue Mackerel East caught by mid-water trawl in the SPF from 2014/15 to 2020/21. n = number of fish. See Table 3-2 for sample N.

# Age structure

#### Mid-water trawl fishery: 2014/15-2020/21

The age structure of Blue Mackerel East caught in mid-water trawls was dominated by fish aged 3 to 4 years in 2014/15 (73%); the age range was 2 to 8 years (age mode: 3 years; Figure 3-4). The modal age was also 3 years in 2015/16 and 2016/17. In 2017/18, the age mode decreased to 2 years (51%), and ages ranged from 1 to 6 years. The modal age remained at 2 years since then, with 95% of the fish ageing  $\geq$ 2 years in 2020/21. The maximum estimated age of Blue Mackerel East from mid-water trawl catch samples was 11 years (Figure 3-4, Table 3-2).



Figure 3-4. Age-frequency distributions of Blue Mackerel East caught by mid-water trawl in the SPF from 2014/15 to 2020/21; n = number of fish. See Table 3-2 for sample N.

# 3.3.2 Blue Mackerel West

# 3.3.2.1 Fishery statistics

#### Number of vessels

The total number of vessels reporting catches of Blue Mackerel West from all jurisdictions declined from a high of 25 vessels in 2008/09 to 10 vessels in 2012/13. Vessel numbers have ranged from 16 to 21 since 2014/15. During 202/21, 18 vessels reported catches of Blue Mackerel West (0 Commonwealth vessels in 2020/21).

#### Annual patterns: Total catch

Total annual catches of Blue Mackerel West were low in the mid/late 1990s and early 2000s (<65 t) and increased to >2000 t in both 2006/07 and 2008/09 (Figure 3-5a). Since then, annual catches decreased to 2 t in 2012/13, remained low through 2014/15 and increased to 983 t in 2015/16 (Figure 3-5a). Total catch decreased from 771 t in 2016/17 to <14 t since then. Historically, the SPF has been the main fishery taking Blue Mackerel West.

#### Annual patterns: Catch, Effort and CPUE

Purse seines have historically been the main gear type used in the SPF for Blue Mackerel West, and purse seining effort has been variable since 2000/01 (Figure 3-5b). Catch and effort peaked in the mid- to late-2000s with effort ranging from 16 to 78 net-sets annually (Figure 3-5b). Purse seine effort and catch in the SPF has decreased since 2007/08, with zero effort reported since 2011/12. Mean annual CPUE of purse seines in the SPF decreased from a peak of 41 t·net-set<sup>-1</sup> in 2007/08 to 9 t·net-set<sup>-1</sup> in 2011/12 (Figure 3-5c).

Prior to 2015/16, mid-water trawl catches of Blue Mackerel West in the SPF were low (Figure 3-5d). Trawl effort increased in the mid-2000s (range: 260–625 trawl-hours) when a multi-purpose 50 m mid-water trawler began targeting small pelagic species, particularly Redbait and Jack Mackerel, off Tasmania. In 2015/16, mid-water trawl effort in the SPF increased to 472 trawl-hours with a Blue Mackerel catch of 979 t (Figure 3-5d). Effort decreased to 365 trawl-hours in 2016/17 (catch: 766 t) and has further decreased to 11.5 trawl-hours in 2017/18 with 0 t of catch. In 2019/20 effort increased to 27 trawl-hours with catch of 9 t; there was no trawl effort in 2020/21. Mean annual CPUE of mid-water trawls in the SPF was <1 t·trawl-hour<sup>-1</sup> prior to 2015/16, 2 t·trawl-hour<sup>-1</sup> in both 2015/16 and 2016/17, and <1 t·trawl-hour<sup>-1</sup> in 2019/20 (Figure 3-5e).



Figure 3-5. Fishery statistics for Blue Mackerel West. (A) Total annual landed catch (tonnes) for all jurisdictions by fishing season from 1984/85 to 2020/21. Long-term trends in the SPF by fishing season from 2000/01 to 2020/21: (B) annual landed catch (tonnes) and effort (net-sets) by purse seine; (C) mean annual CPUE (t·net-set<sup>-1</sup>; ±SE) by purse seine; (D) annual landed catch (tonnes) and effort (trawl-hours) by mid-water trawl; (E) mean annual CPUE (t·trawl-hour<sup>-1</sup>; ±SE) by mid-water trawl.

# 3.3.2.2 Biological Information

Samples of Blue Mackerel West for biological analysis were collected from purse seine catches taken off South Australia from 2008/09 to 2010/11 (Table 3-3). A total of 1,257 fish were sampled over the three years; sex ratios were close to 1:1 but with slightly more females than males (Table 3-3). These catches were limited to the summer/early autumn period, and these biological samples may not be representative of the population. No samples were collected from 2011/12 to 2014/15, due to low levels of fishing activity.

Between 2015/16 and 2020/21, 31 samples were collected in the SPF from midwater trawl catches in the West sub-area (Table 3-3). Length-frequency data were collected from 1,181 Blue Mackerel. Age-frequency data were obtained from 578 of those fish (Table 3-3). Little to no fishing occurred for Blue Mackerel West by the SPF in 2017/18, 2018/19 and 2020/21; no catch sampling occurred in these years.

 Table 3-3. Summary of Blue Mackerel West catch samples collected from commercial SPF landings. Note:

 number of samples in brackets is number for age frequency if different from length frequency.

Season	SPF sub- area	Gear type	No. of samples	Length- frequency n	Age- frequency n	Size range (mm FL)	Age range (years)	Sex ratio M:F
2008/09	West	purse seine	1	79	74	316–390	3–6	1:1
2009/10	West	purse seine	28	933	396	245–400	2–8	0.9:1
2010/11	West	purse seine	8	245	180	293–395	3–8	0.9:1
2015/16	West	mid-water trawl	9	142	102	239–403	0–11	na
2016/17	West	mid-water trawl	21	1,020	464	229–425	2–12	na
2019/20	West	mid-water trawl	1	19	12	221-335	2-4	na

#### Size structure

#### Purse seine fishery: 2008/09–2010/11

Blue Mackerel West sampled during 2008/09 off South Australia ranged from 320 to 390 mm FL (Figure 3-6); >50% of fish were between 340 and 370 mm FL. In 2009/10 and 2010/11, most fish ranged between 300 and 400 mm FL. Annual modal lengths of Blue Mackerel from purse seine catches in the West (350–370 mm FL; Figure 3-6) were larger than those of fish caught in the East (250–310 mm FL; Figures 3-2 and 3-3).

#### Mid-water trawl fishery: 2015/16–2020/21

The modal length of Blue Mackerel West from mid-water trawl catch samples in 2015/16 was 350 mm FL (range: 239–403 mm FL) and increased to 370 mm FL (range: 229–425 mm FL) in 2016/17. Modal length decreased to 300 mm FL in 2019/20 (Figure 3-7; Table 3-3), but the sample size was small. The length-frequency distributions are similar to purse seine catches in the West, and the modal length is larger than in the East (250–310 mm FL; Figures 3-2 and 3-3).



Figure 3-6. Length-frequency distributions of Blue Mackerel West caught by purse seine in the SPF from 2008/09 to 2010/11; n = numberof fish. See Table 3-3 for sample N.



Figure 3-7. Length-frequency distributions of Blue Mackerel West caught by mid-water trawl in the SPF during from 2015/16 to 2019/20; n = number of fish. See Table 3-3 for sample N.

## Age structure

#### Purse seine fishery: 2008/09–2010/11

In 2008/09, the age of fish in purse seine catches ranged from 3 to 6 years (Figure 3-8); 82% of fish were 4–5 years old. A similar age structure was found in 2009/10 and 2010/11, but some fish were up to 8 years old. In all years, most fish were older than 3 years.

#### Mid-water trawl fishery: 2015/16-2020/21

Ages of Blue Mackerel West caught in mid-water trawls ranged from 0 to 12 years (Table 3-3). The age structure shifted towards younger fish in 2019/20 compared to 2015/16 and 2016/17 (Figure 3-9). The modal age in 2015/16 was 5 years and 7 years in 2016/17. In 2019/20, modal age decreased to 3, although sample size was small (Figure 3-9).



Figure 3-8. Age-frequency distribution for Blue Mackerel West caught by purse seine in the SPF from 2008/09 to 2010/11; n = number of fish. See Table 3-3 for sample N.



Figure 3-9. Age-frequency distribution for Blue Mackerel West caught by mid-water trawl in the SPF from 2015/16 to 2019/20; n = number of fish. See Table 3-3 for sample N.

# 3.4 Summary and Conclusions

# 3.4.1 Blue Mackerel East

The main fisheries for Blue Mackerel East are the NSW Ocean Hauling Fishery (purse seine) and the SPF. From 1995/96 to 2014/15, NSW Ocean Fisheries and the SPF took an average of 69% and 20% of the total annual catch in the East sub-area, respectively. Total annual catches from 2010/11 to 2014/15 were <550 t. The SPF has taken >80% of the total catch since 2015/16 and >90% since 2019/20.

The total catch of Blue Mackerel East in the SPF in 2020/21 was 6,202 t, which is the highest on record. Total catches in the SPF increased each year since 2016/17 (1,248 t). The increase in catch reflects the recent establishment of the fishing operation off southern NSW. The catch-per-unit-effort (CPUE) of Blue Mackerel East since 2017/18 of ~4 t·trawl-hour<sup>-1</sup> was similar to that recorded by the factory trawler in 2015/16 (3 t·trawl-hour<sup>-1</sup>) and 2016/17 (4 t·trawl-hour<sup>-1</sup>). The increase in catch of Blue Mackerel East since 2016/17 reflects the increased trawl effort (e.g. 1,483 trawl-hours in 2020/21 compared to 349 trawl-hours in 2016/17).

In 2020/21, the distribution of fish lengths of Blue Mackerel East in mid-water trawl catches was unimodal, with a mode at 250 mm FL. During 2018/19 to 2019/20, the modal length of Blue Mackerel East varied between 240 and 270 mm FL, similar to the mean size of 50% maturity (~260 mm FL). The modal age of Blue Mackerel East from 2018/19 to 2020/21 remained at 2 years. From 2014/15 to 2016/17, the modal size of Blue Mackerel East in mid-water trawl catches was 260 to 280 mm FL while the age mode for those fishing seasons was 3 years. The decrease in the size and age of Blue Mackerel East taken since 2016/17 may reflect changes in the vessels and gear operating in the fishery.

The average total catch of Blue Mackerel East over the last three fishing seasons has been 5,724 t (2020/21: 6,697 t). The spawning biomass of Blue Mackerel for 2019 was 88,265 t (95% CI: 33,320–143,209; Ward et al. 2021a) and is similar to the estimate obtained in 2014 of 83,300 t (Ward et al. 2015b). The recent average exploitation rate has been <10%, which is below the maximum exploitation rate at Tier 1 for Blue Mackerel East of 15% (Smith et al. 2015). The total catch in 2020/21 in the SPF was 52% of the available TAC. On the basis of the information provided here, Blue Mackerel East is classified as sustainable. Piddocke et al. (2021) also found the biological stock of Blue Mackerel East to be sustainable. Patterson et al. (2021) classified Blue Mackerel East as 'not overfished' and 'not subject to overfishing'.

# 3.4.2 Blue Mackerel West

Total annual catches of Blue Mackerel West were low in the late 1990s and early 2000s (<55 t) and increased to >2000 t in 2006/07 and 2008/09. In the SPF, catches have been mainly taken by purse seining prior to 2015/16. Total annual catches decreased to <2 t from 2012/13-2014/15, increased to 979 t in 2015/16, and decreased to 766 t in 2016/17.

There was negligible catch of Blue Mackerel West between 2017/18 and 2020/21. Mean annual CPUE of purse seines in the SPF peaked at 41 t·net-set<sup>-1</sup> in 2007/08; CPUE of mid-water trawls rose to 2 t·trawl-hour<sup>-1</sup> in 2015/16 and 2016/17 and has been <1% since then. Low annual catch in recent years reflects low fishing effort in areas where Blue Mackerel are known to be abundant (e.g. Great Australian Bight).

Blue Mackerel from commercial catch samples from both purse seines and mid-water trawls in the West have been well above the mean size at 50% maturity of ~260 mm FL (Ward and Rogers 2007) and tend to be larger and older than those from the East. In 2015/16, the modal length of Blue Mackerel West from mid-water trawls was 350 mm FL and increased to 370 mm FL in 2016/17. There was a decrease in modal length to 300 mm in 2019/20, but the sample size was small. The modal age class increased from 5 years in 2015/16 to 7 years in 2016/17 (age range: 0–12 years). Modal length decreased to 3 years in 2019/20 (1 sample, 12 fish). Age and length structures are difficult to interpret due to the limited fishing effort, small sample sizes, differing ageing methods and changes in fishing locations over time.

A preliminary application of the DEPM to Blue Mackerel West off South Australia during 2005 provided a 'best' estimate spawning biomass of 56,228 t (Ward and Rogers 2007). This estimate of spawning biomass was considered to be conservative because the survey only covered a limited part of the West sub-area, and there was clear evidence of significant spawning activity outside the survey area in the western Great Australian Bight (Ward and Rogers 2007).

Recent catches of Blue Mackerel West have been <1% of the estimated spawning biomass for 2005 and below the Tier 3 exploitation rate for this stock of 3.75%. In 2019/20, the annual catch in the SPF (9 t) was <1% of the available TAC. On the basis of the information provided here, Blue Mackerel West is classified as sustainable. Piddocke et al. (2021) classified the biological stock of Blue Mackerel West as sustainable. Patterson et al. (2021) classified Blue Mackerel West as 'not overfished' and 'not subject to overfishing'.

# 4 Redbait (*Emmelichthys nitidus*)

# 4.1 Introduction

# 4.1.1 Background to Fishery

Redbait was a key by-product species in the Tasmanian Purse Seine Fishery for Jack Mackerel (*Trachurus declivis*) that developed off Tasmania in the mid-1980s. This fishery has recorded catch and effort data since its inception in 1984. Logbooks contain a record of fishing operations and species taken by net-set. Although landings of Redbait rarely exceeded 5% of the total catch in a net-set, annual catches averaged ~700 t from 1984/85 to 1989/90 (Pullen 1994).

Mid-water trawling to target subsurface schools of Jack Mackerel off Tasmania was trialled in 2001/02 (Welsford and Lyle 2003). Between December 2001 and April 2002, a total catch of over 5,000 t of small pelagic fishes was taken; 90% was Redbait. In late 2002, a multi-purpose 50 m mid-water trawler began targeting small pelagic species off Tasmania, particularly Redbait. By mid-2003, more than 7,000 t of small pelagic fishes had been taken, with Redbait dominating the catch. Small-scale purse seine operations were temporarily resumed in response to declining trawl effort in the late 2000s (Emery et al. 2015).

Redbait have primarily been frozen whole for use as feed for farmed Southern Bluefin Tuna (*Thunnus maccoyii*) and have also been used to produce fish meal for the aquaculture industry.

# 4.1.2 Taxonomy

Redbait (*Emmelichthys nitidus,* Richardson 1845) belong to the family *Emmelichthyidae*, which contains three genera and 15 species (Nelson 2006). Redbait are one of two species of emmelichthyid found off southern Australia, the other being the Rubyfish (*Plagiogeneion rubiginosum*) (Last et al. 1983, May and Maxwell 1986, Gomon et al. 2008).

# 4.1.3 Distribution

Emmelichthyids are found throughout tropical and temperate waters world-wide. Generally, they are found in schools over continental shelf breaks, seamounts and submarine ridges. They inhabit depths from the surface to >800 m, though are mostly recorded from mid-water trawls in 100–400 m water (Heemstra and Randall 1977, Smith and Heemstra 1986, Mel'nikov and Ivanin 1995). Redbait are widely distributed throughout the southern hemisphere, with the species reported from Tristan da Cunha in the southern Atlantic, the south-western coast of South Africa, St Paul and Amsterdam Islands, midoceanic ridges and seamounts through the Indian Ocean, Australia, New Zealand, submarine ridges in the south-eastern Pacific, and the southern coast of Chile (Markina and Boldyrev 1980, Meléndez and Céspedes 1986, Parin et al. 1997). Within Australian waters, their range extends from mid NSW to south-west Western Australia, including Tasmania (Gomon et al. 2008).

# 4.1.4 Stock Structure

There have been no studies on the stock structure of Redbait in Australia. However, Redbait from eastern Australia are thought to be a single stock based on spawning dynamics (Bulman et al. 2008). The situation for western Tasmania and the GAB is less clear. Neira et al. (2008) observed biological differences in Redbait from eastern and south-western Tasmania, but those differences may have resulted from sampling bias due to the different depths fished in each region (Ewing and Lyle 2009).

Egg distributions from spawning biomass surveys indicate Redbait in south-eastern Australia may be a continuous stock along the continental shelf break and inner slope from southern NSW, around Tasmania to Kangaroo Island in South Australia (Neira et al. 2008, Ward et al. 2019, Grammer et al. 2022). There was no evidence of spawning activity through Bass Strait (Ward et al. 2019, Grammer et al. 2022).

## 4.1.5 Movement

No studies have investigated movement of Redbait.

# 4.1.6 Food and Feeding

In South African coastal waters, smaller size classes of Redbait (136-280 mm) feed exclusively on small planktonic crustaceans, with euphausiids (*Nyctiphanes* and *Euphausia* spp.), hyperiid amphipods (primarily *Themisto gaudichaudi*), mysids and large copepods comprising the entire diet (Meyer and Smale 1991). Larger Redbait (281–493 mm) also fed primarily on small planktonic crustaceans, but nekton, such as cephalopods, carid shrimp, and small fishes including myctophids, were part of the diet (Meyer and Smale 1991). Redbait captured on the shelf off eastern Victoria (unspecified size) had a varied diet that was dominated by pelagic crustaceans and other invertebrates, including gelatinous zooplankton (Bulman et al. 2000, Bulman et al. 2001). Similarly, Redbait captured off eastern Tasmania consumed mainly pelagic crustaceans, with krill and copepods comprising 66% and 33% of the diet, respectively (McLeod et al. 2012).

The diet of Redbait is similar to that of Jack Mackerel from Tasmania, with krill representing the dominant prey item on the continental shelf (Young et al. 1993, McLeod et al. 2012). Since Redbait and Jack Mackerel form mixed species schools in Tasmanian waters (Williams and Pullen 1993), it is not surprising the two species feed on similar prey.

# 4.1.7 Age, Growth and Size

The maximum reported size for female and male Redbait from Tasmania is 317 and 304 mm FL, respectively (Neira et al. 2008), which is considerably smaller than reported in other areas. Redbait grow to 335 mm FL off eastern Victoria (Furlani et al. 2000), 344 mm standard length (SL) off the coast of Chile (Meléndez and Céspedes 1986) and to 493 mm TL in South African waters (Heemstra and Randall 1977, Meyer and Smale 1991). Redbait are observed to school by size and stratify by water depth, with larger (>200 mm FL) individuals found deeper and closer to the seafloor (Markina and Boldyrev 1980).

Growth estimates for Redbait (otolith-based) suggest rapid growth during the first few years (Williams et al. 1987, Neira et al. 2008). On average, Redbait off Tasmania reached >200 mm FL in the first three years, with growth slowing thereafter (Neira et al. 2008). Differences in growth rates have been reported for Redbait off Kangaroo Island compared to those off southern NSW; fish off Kangaroo Island were larger at a given age than those off southern NSW (Dennis et al. 2021).

The maximum estimated age for Redbait is 21 years for females and 18 years for males (Neira et al. 2008). The larger Redbait reported from Africa (e.g. Meyer and Smale 1991) suggest that maximum age may be higher than reported for Tasmanian fish or that growth rates vary between regions. Age validation of Rubyfish in New Zealand, using otoliths and the bomb radiocarbon chronometer, has shown fish over 400 mm can be up to 100 years old (Paul et al. 2000, Horn et al. 2012), indicating some emmelichthyids are long-lived.

#### 4.1.8 Reproduction

Redbait is an asynchronous batch spawner with indeterminate fecundity (Neira et al. 2008, Ward et al. 2019). Annual trends in Gonadosomatic Index (GSI) and macroscopic gonad stages indicated that peak spawning season of Redbait occurs off eastern Tasmania during September and October, and off south-western Tasmania during October (Ewing and Lyle 2009). Most spawning occurs along the continental shelf break and inner slope where water depths are approximately 200 m and water temperatures are between 12 and 16°C (Neira et al. 2008, Ward et al. 2019, Grammer et al. 2022).

Regional differences in size and age at sexual maturity for Redbait have been noted, but Ewing and Lyle (2009) suggested this difference could have resulted from sampling bias due to the different depths fished in each region. Males and females from south-western Tasmania matured ~100 mm larger and two years older than Redbait from eastern Tasmania (Ewing and Lyle 2009). The size (age) at 50% sexual maturity for Redbait in eastern Tasmania was 147 mm FL (2 years) for males and 157 mm FL (2 years) for females. In south-western Tasmania, the size (age) at 50% sexual maturity was 244 mm FL (4.8 years) for males and 261 mm FL (4.1 years) for females (Ewing and Lyle 2009).

During recent DEPM surveys for Redbait, relative fecundity has been estimated to be ~85 oocytes per gram of weight for fish off south-eastern Australia (Ward et al. 2019, Grammer et al. 2022). Mean spawning fraction (proportion of mature females spawning per

Securing Australia's fishing future

day/night) is estimated to be ~0.200 in south-eastern Australia (Ward et al. 2019, Grammer et al. 2022).

# 4.1.9 Early Life History and Recruitment

Redbait eggs are positively buoyant and hatch about 2.5–4.5 days after fertilisation depending on temperature (e.g. 13.1–16.5°C; Neira et al. 2008). Newly hatched yolk sac larvae range from 1.9–3.3 mm TL. Little is known about the early life history of Redbait. The distribution of eggs and larvae off south-eastern Australia have been described by Neira et al. (2008), Ward et al. (2019) and Grammer et al. (2022).

## 4.1.10 Stock Assessment

The first dedicated DEPM survey for Redbait East was conducted in 2020 and extended from southern NSW to southern Tasmania (Grammer et al. 2022). A preliminary DEPM survey was done in 2005 and 2006 that covered the area from the north-eastern Bass Strait to south of the Tasman Peninsula, but didn't include the entire spawning area (Neira et al. 2008; Ewing and Lyle 2009; Neira and Lyle 2011). The first dedicated DEPM survey for Redbait West along the southern Australian coast (Kangaroo Island to western Tasmania) occurred during October 2017 (Ward et al. 2019).

# 4.1.11 Recreational fishing

There is no known recreational fishery for Redbait in Australia.

# 4.1.12 Biomass Estimates

#### East

The DEPM was applied to Redbait East during 2005 and 2006 (Neira et al. 2008) and in 2020 (Grammer et al. 2022). The preliminary application of the DEPM estimated the spawning biomass of Redbait East from the north-eastern Bass Strait to south of the Tasman Peninsula during October 2005 and 2006 to be 86,994 t and 50,782 t, respectively (Neira and Lyle 2011). These estimates are considered to be negatively biased, as the surveys covered less than half the known spawning area of Redbait in south-eastern Australia (Neira and Lyle 2011).

The first dedicated application of the DEPM to Rebait East was undertaken in October 2020 off south-eastern Australia; the estimate of spawning biomass for the survey was 52,629 t (95% CI = 13,937–91,321) (Grammer et al. 2022). The estimated spawning area off south-eastern Australia was 19,715 km<sup>2</sup>, comprising 34.2 % of the total area sampled (57,486 km<sup>2</sup>) (Grammer et al. 2022). Live Redbait eggs (n = 2,372) were collected from 79 of the 231 (34.2%) stations between Jervis Bay, NSW and South East Cape, Tasmania. Mean daily egg production ( $P_0$ ) was 22.7 eggs·day<sup>-1</sup>·m<sup>-2</sup>. The highest densities of Redbait eggs were recorded in mid to outer shelf waters between northern Bass Strait and southern NSW; SSTs ranged between 11 and 16°C.

Sensitivity analyses showed that realistic variations of each parameter produced estimates of spawning biomass for Redbait East that were between about 50,000 t and 75,000 t. Variations in the values of  $P_0$  caused the greatest change in estimates of spawning biomass (Grammer et al. 2022). The estimate of spawning biomass for Redbait East is considered to be reliable, because it is based on robust and/or conservative estimates of all key parameters.

# West

The first dedicated application of the DEPM to Rebait West was undertaken in October 2017 between Kangaroo Island, South Australia and south-western Tasmania; the estimate of spawning biomass for the survey was 66,767 t (CI = 28,797–190,392) (Ward et al. 2019). The estimated spawning area off southern Australia was 28,365 km<sup>2</sup>, comprising 36.3 % of the total area sampled (78,212 km<sup>2</sup>) (Ward et al. 2019). Live Redbait eggs (n = 3,280) were collected from 113 of the 308 (36.7%) stations. Mean daily egg production ( $P_0$ ) was 22.5 eggs·day<sup>-1</sup>·m<sup>-2</sup>. The highest densities of Redbait eggs were recorded in outer shelf waters off western Victoria and the west coast of Tasmania; SSTs ranged between 12 and 15°C.

The estimate of spawning biomass for Redbait West is considered reliable, because it is based on robust and/or conservative estimate of key parameters. This estimate is also similar to the Atlantis-SPF biomass estimate for Redbait West of 66,000 t (typical range: 59,000–70,000 t; Smith et al. 2015).

# 4.1.13 Management Strategy Evaluation

Smith et al. (2015) concluded the harvest rate of 15% may be too high for Redbait and suggested a Tier 1 harvest rate of 9% for Redbait East and 10% for Redbait West, with the Tier 1 rate being applied for not more than 5 years. Tier 2 harvest rates for Redbait East and West were recommended to be 50% of Tier 1 rates and not to be applied for more than 10 years. The study also indicated that it is not safe to apply Tier 2 harvest rates unchecked for long periods of time (Smith et al. 2015). The Atlantis-SPF biomass estimate for Redbait East is 82,000 t (typical range: 75,000–105,000 t) and 66,000 t (typical range: 59,000–70,000 t) for Redbait West (Smith et al. 2015).

#### 4.1.14 Management

Currently, Redbait East is managed at the Tier 2 level (transitioning to Tier 1 in the 2022/23 fishing season) and Redbait West is managed at the Tier 1 level under the Harvest Strategy. DEPM assessments of Redbait have been conducted for both the East and West sub-areas of the SPF: Redbait East in 2005, 2006, and 2020, and Redbait West in 2017.

# 4.2 Methods

# 4.2.1 Fishery Statistics

Fishery statistics from 1995/96 to 2020/21 were provided by relevant jurisdictions and collated by SARDI Aquatic Sciences. Annual data are reported in fishing seasons (May 1 to April 30) rather than financial years as was done in previous assessments (e.g. Ward et al. 2013, 2014c, 2015c).

Estimates of total annual catch supplied for Redbait East include data from the NSW Ocean Fisheries (Hauling, Trap and Line, Trawl), Victorian Ocean Purse Seine Fishery, Tasmanian Scalefish Fishery and the Commonwealth SPF. In the West, total annual catch estimates include data from the Tasmanian Scalefish Fishery and Commonwealth SPF. Due to data confidentiality (<6 licence holders annually reporting catch since 2015/16), fishery data from Victoria were not provided and have not been included in total annual catch statistics since 2015/16.

Mean annual CPUE of Redbait in the Commonwealth SPF by sub-area is calculated for the gear types of mid-water trawl (tonnes·trawl-hour<sup>-1</sup> ±SE) and purse-seine (tonnes·net-set<sup>-1</sup> ±SE) from 2000/01 to 2020/21. For the 2020/21 fishing season, shots by mid-water trawl of <0.2 hrs were removed from the analyses, as these were deemed 'false shots' (pers. comm. AFMA). Zero catch of Redbait in a trawl was assumed when effort but not catch was reported in the logbook record.

#### 4.2.2 Biological Information

Fishery-dependent length frequency and biological data were collected for Redbait between 1984 and 1993 as part of a monitoring program for the Jack Mackerel Purse seine Fishery off Tasmania. Some biological information was obtained from samples collected between 1985 and 1990 from demersal research trawls conducted by CSIRO and the Tasmanian fisheries agency. Between 1994 and 2001, there was limited catch sampling of the purse seine fishery.

Biological data were collected by AFMA observers from a small proportion of trips during the 2001/02 pair-trawl fishing trials undertaken off Tasmania. When mid-water trawl operations started in 2002, TAFI began an intensive biological monitoring program that continued to 2006. AFMA also provided observer coverage of mid-water trawl operations, with additional length-frequency data collected from 2002 to 2008.

Purse seine operations for small pelagic fish resumed in Tasmanian State waters in 2008/09, mainly targeting Redbait and Jack Mackerel. Catch sampling of mid-water trawl and purse seine operations adjacent to Tasmania began in 2009/10 as part of the SPF monitoring program under the Harvest Strategy (AFMA 2008). Catch samples were not collected for Redbait from 2010/11 to 2013/14 due to limited fishing activity. Catch sampling by AFMA observers resumed in the SPF during 2014/15. Samples of Redbait

were collected (n = 50 randomly selected fish per trawl) and supplied to SARDI Aquatic Sciences to estimate the size and age composition of the catch.

Biological data collected from each fish include: body length (mm FL), total weight (±1 g), sex, gonad developmental stage (following the macroscopic staging criteria described in Neira et al. 2008) and gonad weight (±0.1 g). Gonad stages were designated as: I) immature; II) maturing virgins or recovering spent; III) maturing; IV) ripe; and V) spent. Otoliths were removed from random sub-samples of the fish for age determination. The age structure of Redbait prior to 2014/15 was estimated using age-length keys based on age data pooled between 2001/02 to 2005/06. From 2014/15 to present, ages for Redbait have been based on annual growth increment counts in thin-sectioned otoliths (sub-samples of 5 to 10 fish per sample).

Redbait otoliths were aged following the protocol of Ewing and Lyle (2009) after thinsectioning and mounting on microscope slides. Edges of each annual opaque zone were counted (edges of opaque zones are more easily delineated than edges of translucent zones). Identifying the edge of the first annual opaque zone followed guidelines described in (Ewing and Lyle 2009), where it followed a distinctive wide translucent area just exterior to the inner opaque region. The period of age represented by the opaque core was assumed to be similar across all regions where the fish were collected based on spawning stock, spawning time and time of opaque increment formation (Neira et al. 2008, Ewing and Lyle 2009). All fish were assigned a birthdate of Oct 1 based on the spawning times for the spawning stocks off southern Australia (Ewing and Lyle 2009). Redbait form an opaque zone during winter with the completion becoming discernible from November to February (Ewing and Lyle 2009). The completion of annual opaque zones in otoliths during spring/summer in temperate marine waters is common (e.g. Choat and Axe 1996, Fowler and Short 1998, Smith and Deguara 2003, Ewing et al. 2007). Therefore, we designated a standardised completion date of Oct 31 for the opaque zone for all regions. The first year of growth of Redbait in southern Australia represents growth from 1 Oct of the birth year to 31 Oct of the next year (13 months). Each otolith was aged by two different readers and the counts compared. Where the counts differed, the otolith was re-read and an age consensus reach, or the otolith was removed from further analyses. Prior to ageing otoliths, the reader used a species specific reference collection to calibrate their ageing technique (see Ward and Grammer 2019).

Catch weighting was applied to length- and age-frequency data collected since 2014/15 in each sub-area. Length- and age-frequencies were weighted by the number of fish sampled per trawl to account for uneven sample sizes and then were catch weighted by the total amount of Redbait taken in the same trawl.

Commercial logbook information, length-frequency and biological data collected between 1984 and 2021 are included in this assessment. In addition to current catch samples, age, growth and reproductive data for Redbait were available from previous studies (i.e. Welsford and Lyle 2003 and Neira et al. 2008). Summarised biological data prior to

2014/15 are presented in financial years. From 2014/15 to present, all SPF catch sampling data are presented in fishing seasons from 1 May to 30 April.

# 4.3 Results

## 4.3.1 Redbait East

# 4.3.1.1 Fishery Statistics

#### Number of vessels

Since 1995/96, a limited number of vessels from all jurisdictions have reported catches of Redbait East; total annual vessel numbers have ranged from zero to seven. On average, over the last 25 years, three boats per year have reported catches of Redbait East, and ~40% of the vessels reporting catch in each year are Commonwealth vessels. From 2013/14 to 2017/18 and in 2019/20, 1 vessel annually landed Redbait East; the number vessels increased to 3 in 2018/19 and 2020/21.

#### Annual patterns: Total catch

Due to data confidentiality (<6 licence holders reporting catch per year), only SPF catches are reported in some years. From 1995/96 to 2000/01, Redbait East catches did not exceed 315 t, and purse seining was the prevailing method (Figure 4-1a). Mid-water trawling began to replace purse seining in the early 2000s (Figure 4-1a). With the change to mid-water trawling, catches increased to 3,610 t in 2001/02 and peaked at 7,728 t in 2003/04. From then onwards, annual catches declined to 75 t in 2010/11. No catches of Redbait East were reported from 2011/12 to 2013/14; 217 t were taken in 2015/16 and catches declined to 9.5 t in 2017/18. Catches increased to 539 t in 2018/19, 2,457 t in 2019/20, and decreased to 2,011 t in 2020/21.

#### Annual patterns: Catch, Effort and CPUE

There has been limited use of purse seines in the SPF for Redbait East; mid-water trawls have historically been the main gear type (Figures 4-1a). Since 2000/01, purse seine effort has not exceeded 7 net-sets with a maximum catch of 54 t in 2004/05 (Figure 4-1b). There has been no reported purse seine effort and catch of Redbait East in the SPF since 2007/08 (Figure 4-1b). Mean annual CPUE of purse seining in the East peaked in 2004/05 (8 t·net-set<sup>-1</sup>) and decreased to ≤3 t·net-set<sup>-1</sup> in 2005/06–2007/08, with minimal to no fishing in other years (Figure 4-1c).

Mid-water trawl effort in the SPF for Redbait East reached 1,338 trawl-hours (catch: 7,721 t) in 2003/04, decreased to 0 trawl-hours (2011/12 to 2013/14) and increased to 638 trawl-hours (catch: 217 t) in 2015/16 (Figure 4-1d). In 2016/17 effort decreased to 349 trawl-hours (catch: 101 t), and increased to 1,307 in 2019/20 (catch: 2457 t) and 1,483 trawl-hours in 2020/21 (catch: 2011 t).

Mean annual CPUE of Redbait in mid-water trawls in the SPF declined from 8 t·trawl-hour<sup>1</sup> in 2001/02 to 1 t·trawl-hour<sup>1</sup> in 2010/11, with no further activity until 2014/15 (Figure 4-1e). Mean annual CPUE was <1 t·trawl-hour<sup>1</sup> from 2015/16 to 2018/19, increasing to <2 t·trawl-hour<sup>1</sup> in 2019/20 and 2020/21 (Figure 4-1e).



Figure 4-1. Fishery statistics for Redbait East. (A) Total annual landed catch (tonnes) for all jurisdictions from 1984/85 to 2020/21. Black bars: catch per financial year; grey bars: catch per fishing season. ( $\star$ ) indicates data confidentiality and only catches by the SPF are shown. Long-term trends in the SPF by fishing season from 2000/01 to 2020/21: (B) annual landed catch (tonnes) and effort (net-sets) by purse seine; (C) mean annual CPUE (t-net-set<sup>-1</sup>; ±SE) by purse seine; (D) annual landed catch (tonnes) and effort (trawl-hours) by mid-water trawl; (E) mean annual CPUE (t-trawl-hour<sup>-1</sup>; ±SE) by mid-water trawl.

# 4.3.1.2 Biological Information

Length-frequency data were collected from 36,804 Redbait sampled from commercial midwater trawl catches off south-western Tasmania between 2001/02 and 2009/10 (Table 4-1). Age-frequency data were obtained from 35,980 of those fish (Table 4-1). Catch samples were not collected from 2010/11 to 2014/15 due to low levels of fishing activity in the East sub-area. Between 2014/15 and 2010/21, 102 samples of Redbait were collected in the SPF from midwater trawl catches in the East sub-area (Table 4-1). Length-frequency data were collected from 3,913 Redbait. Age-frequency data were obtained from 1,139 of those fish (Table 4-1).

Season	SPF sub- area	Gear type	No. of samples	Length- frequency n	Age- frequency n	Size range (mm FL)	Age range (years)
2001/02	East	mid-water trawl	-	1,523	1,523	110-290	1-19
2002/03	East	mid-water trawl	-	5,036	5,036	90-290	1-19
2003/04	East	mid-water trawl	-	13,126	13,126	110-290	1-19
2004/05	East	mid-water trawl	-	12,225	12,225	100-300	1-19
2005/06	East	mid-water trawl	-	5,473	5,473	120-290	1-19
2007/08	East	mid-water trawl	-	551	-	100-260	-
2009/10	East	mid-water trawl	6	393	120	170–300	1–9
2014/15	East	mid-water trawl	2	61	40	200–275	2–14
2015/16	East	mid-water trawl	35	2,091	439	87–277	0–15
2016/17	East	mid-water trawl	5	242	56	119–278	2–14
2017/18	East	mid-water trawl	3	86	29	210–261	2–15
2018/19	East	mid-water trawl	14	636	165	151-224	1-6
2019/20	East	mid-water trawl	23	536	318	150-234	1-3
2020/21	East	mid-water trawl	20	261	92	118-220	0-3

#### Table 4-1. Summary of Redbait East catch samples collected from commercial SPF landings.

#### Size structure

#### The purse seine fishery: 1984/85 to 1993/94 and 2009/10

Purse seine catches of Redbait between 1984/85 and 1994/95 off eastern Tasmania mainly contained fish between 140 and 290 mm FL (max length: 320 mm FL; Figure 4-2). Catches between 1984/85 and 1987/88 were dominated by fish from 200–300 mm FL, with only a few small fish (100–140 mm FL) caught in 1985/86. A strong cohort of smaller fish (120–170 mm FL) was present in the size structure for 1988/89 and accounted for most of the catch in the following year. Between 1989/90 and 2009/10, smaller fish (160–

240 mm FL) were prevalent (Figure 4-2). No catch samples have been collected from purse seines since 2009/10.



Figure 4-2. Length-frequency distributions (mm FL) of Redbait caught in the SPF by purse seine from 1984/85 to 1993/94 and in 2009/10. n = number of fish.

#### Mid-water trawl fishery: 2001/02–2020/21

Redbait East caught by mid-water trawl operations between 2001/02 and 2007/08 off eastern Tasmania were considerably smaller than those caught by the earlier purse seine operations, with individuals mainly between 100 and 200 mm FL (Figures 4-2 and 4-3). Redbait East catches contained a high proportion of small fish with modes varying between 110 and 180 mm FL (Figure 4-3). Only a small proportion of the catch was made up of fish larger than 200 mm FL. The size structure of Redbait East in mid-water trawl catches increased during 2009/10; fish were primarily 190–240 mm FL (modal length: 190 mm FL; Figure 4-3).

From 2014/15 to 2017/18, modal lengths were larger than those from prior years and varied between 200 to 240 mm FL. Modal length of Redbait East dropped to 170 mm in 2018/19 and 2019/20 and increased to 180 mm in 2020/21. (Figures 4-3, Table 4-1).



Figure 4-3. Length-frequency distributions of Redbait East caught by mid-water trawl in the SPF from 2001/02 to 2020/21. n = number of fish. See Table 4-1 for sample N since 2009/10.

## Age structure

#### The purse seine fishery: 2009/10

Purse seine catches off eastern Tasmania mainly contained 2–3 year old fish in 2009/10, with 2 year olds making up >50% of the catch (Figure 4-4). Age data for Redbait East caught by purse seine in the SPF were not available for other years.



Figure 4-4. Age-frequency distribution of Redbait East caught by purse seine in the SPF during 2009/10; n = number of fish.

#### Mid-water trawl fishery: 2001/02–2020/21

Mid-water trawl catches of Redbait East prior to 2014/15 mainly contained fish between 1 and 5 years with maximum ages of 19 years (Figure 4-5, Table 4-1). Catches off eastern Tasmania contained mostly fish aged 1 and 2 years from 2001/02 to 2002/03. During 2003/04 to 2009/10, the age structure shifted to fish aged 2 and 3 years (Figure 4-5). Mid-water trawl catches of 2009/10 had slightly older age structures than purse seine catches in the same year (Table 4-1; Figures 4.4–4.5).

In 2014/15, the age structure had an increasing proportion of older fish: 57% of the fish were >4 years old (Figure 4-5). Two modes are present in the age structure during 2015/16: one at 2 years and one at 7 years. Older ages are present in 2016/17, but fish aged 2 years dominate the catch. The age structure in 2017/18 was based on a small number of samples (N = 3, n = 29) and shows both young and old fish were prevalent. In the 2018/19 to 2020/21 fishing seasons, >80% of the fish were 2 years or less (Figure 4-5). Since 2014/15, the maximum age found in mid-water trawl catches has been 15 years.



Figure 4-5. Age-frequency distributions of Redbait East caught by mid-water trawl in the SPF from 2001/02 to 2020/21; n = number of fish. See Table 4-1 for sample N since 2009/10.

# 4.3.2 Redbait West

# 4.3.2.1 Fishery Statistics

#### Number of vessels

In the West sub-area of the SPF, the number of vessels reporting catches of Redbait since 1995/96 were lower than those in the East: the annual average has been 1 boat per year. The SPF is the principal fishery reporting catches of Redbait West. From 2011/12 to 2013/14, no vessels reported catches of Redbait West. From 2014/15 to 2017/18, 1 vessel annually has fished for Redbait in the West sub-area. In 2018/19 and 2020/21, no vessels reported catch of Redbait West; 1 vessel reported catches in 2019/20.

#### Annual patterns: Total catch

Due to data confidentiality (<6 licence holders reporting catch per year), only SPF catches are reported in some years. Historically, Redbait catches in the West sub-area have been lower than those in the East, but follow a similar temporal trend (Figure 4-6a). Mid-water trawling has been the primary method used to target Redbait West. From 1995/96 to 2000/01, there were minimal catches of Redbait West. Annual catches began increasing in 2001/02, peaked at 3,228 t in 2006/07 and declined to 298 t in 2009/10 (Figure 4-6a). Redbait was not caught again in the West sub-area until 2015/16 (1,157 t). Minimal to no fishing has occurred since 2016/17; 9 t were reported in 2019/20 (Figure 4-6a).

#### Annual patterns: Catch, Effort and CPUE

In the SPF during the mid-2000s, mid-water trawling effort for Redbait West (442–625 trawl-hours) peaked in years when the catch was highest (2,511–3,228 t; Figure 4-6b). No fishing effort was reported for Redbait West from 2010/11 to 2013/14 (Figure 4-6b). Effort began increasing in 2014/15, peaked at 472 trawl-hours (catch: 1,157 t) in 2015/16, and decreased to 11.5 trawl-hours (catch: 0 t) in 2017/18 (Figure 4-6b). Effort decreased to 0 trawl-hours in 2018/19, increased to 27 trawl-hours in 2019/20 (catch: 9 t), and dropped to 0 trawl-hours in 2020/21. Mean annual CPUE of mid-water trawls for Redbait West decreased from 11 t-trawl-hour<sup>1</sup> in 2001/02 to 4 t-trawl-hour<sup>1</sup> in 2009/10, with no effort reported until 2014/15 (Figure 4-6c). Mean annual CPUE increased to 3 t-trawl-hour<sup>1</sup> in 2016/17, dropped to 0 t-trawl-hour<sup>1</sup> in 2017/18 but increased to <1 t-trawl-hour<sup>1</sup> in 2019/20 (Figure 4-6c).



Figure 4-6. Fishery statistics for Redbait West. (A) Total annual landed catch (tonnes) for all jurisdictions from 1995/96 to 2020/21. (\*) indicates data confidentiality and only catches by the SPF are shown. Long-term trends of mid-water trawling in the SPF by fishing season from 2000/01 to 2020/21: (B) annual landed catch (tonnes) and effort (trawl-hours); (C) mean annual CPUE (t·trawl-hour<sup>-1</sup>; ±SE).

# 4.3.2.2 Biological Information

Length-frequency data were collected from 11,797 Redbait sampled from commercial midwater trawl catches off south-western Tasmania between 2001/02 and 2009/10 (Table 4-2). Age-frequency data were obtained from 11,096 of those fish (Table 4-2). Catch samples were not collected from 2010/11 to 2014/15 and in 2020/21 due to low levels of fishing activity in the West sub-area. Between 2015/16 and 2019/20, 45 samples of Redbait were collected in the SPF from midwater trawl catches in the West sub-area (Table 4-2). Length-frequency data were collected from 2,736 Redbait. Age-frequency data were obtained from 651 of those fish (Table 4-2).

Season	SPF sub- area	Gear type	No. of samples	Length- frequency n	Age- frequency n	Size range (mm FL)	Age range (years)
2001/02	West	mid-water trawl	-	150	-	120-170	-
2002/03	West	mid-water trawl	-	1,166	1,166	160-300	2-18
2003/04	West	mid-water trawl	-	574	572	120-290	2-18
2004/05	West	mid-water trawl	-	4,126	4,124	120-300	2-18
2005/06	West	mid-water trawl	-	5,219	5,214	60-310	2-18
2006/07	West	mid-water trawl	-	485	-	170-320	-
2009/10	West	mid-water trawl	1	77	20	210–310	2–13
2015/16	West	mid-water trawl	19	1,526	380	94–291	0–10
2016/17	West	mid-water trawl	25	1,190	259	148–304	1–14
2019/20	West	mid-water trawl	1	20	12	120-200	1-2

Table 4-2. Summary of Redbait West catch samples collected from commercial SPF landings.

# Size structure

# Mid-water trawl fishery: 2001/02–2019/20

Redbait West from mid-water trawl catches off south-western Tasmania during 2001/02 to 2007/08 ranged mainly from 130–280 mm FL (overall modal length: 200 mm FL; Figure 4-7). Redbait West taken over this time period were larger than those from the East (overall modal length: 160 mm FL; Figure 4-3). A single catch of Redbait West in 2009/10 (from south-western Tasmania) contained fish between 210–310 mm FL (mode 240 mm) (Table 4-2). The size structure was bimodal in 2015/16 with modes at 130 and 220 mm FL. In 2016/17, the modal length increased to 240 mm FL. Modal length decreased to 190 mm FL in 2019/20, but sample size was small. The size structure of Redbait West in 2016/17 is similar to those of the purse seine catches in the mid-1980s off eastern Tasmania (Figures 4-2 and 4-7).


Figure 4-7. Length-frequency distributions of Redbait West caught by mid-water trawl in the SPF from 2001/02 to 2019/20. n = number of fish. See Table 4-2 for sample N.

#### Age structure

#### Mid-water trawl fishery: 2002/03-2019/20

From 2002/03 to 2007/08, catches of Redbait West off south-west Tasmania had a higher proportion of older fish than in the East; Redbait West ages ranged from 2–18 years (Figure 4-8, Table 4-2). There was a strong cohort of age 2 fish in 2003/04 catches that was prevalent in 2004/05 catches as 3 years olds and 4 year olds in 2005/06 (Figure 4-8). One sample of Redbait West from south-western Tasmania in 2009/10 had fish that were older than those from 2009/10 catches in the East (up to 13 years; Table 4-2), with 90% of

fish estimated to be over 4 years of age (East: 76%  $\leq$ 4 years). In 2015/16, the age structure ranged from 0–10 years, with 88% aged 2 to 5 years and a modal age of 4 years. The modal age decreased to 3 years in 2016/17, but with an age range of 1 to 14 years. Modal age decreased to 1 year in 2019/20, but sample size was small (Figure 4-8; Table 4-2).



Figure 4-8. Age-frequency distributions of Redbait West caught by mid-water trawl in the SPF from 2002/03 to 2019/20; n = number of fish. See Table 4-2 for sample N.

# 4.4 Summary and Conclusions

## 4.4.1 Redbait East

The main fishery for Redbait East is the SPF. The highest annual catch of Redbait East in the SPF between 2010/11 and 2020/21 was 2,457 t in 2019/20. The annual catch in the SPF in 2020/21 was 2,011 t. Catches of Redbait East in mid-water trawls have mainly contained fish above the mean size at 50% maturity of ~150 mm FL. The spawning biomass of Redbait East was estimated to be ~70,000 t in 2005 and 2006 (Neira et al. 2008). The DEPM was applied to Redbait East in 2020 and provided an estimate of spawning biomass of 52,629 t (95% CI = 13,937–91,321) (Grammer et al. 2022). Redbait East will return to Tier 1 for the 2022/23 fishing season with a maximum exploitation rate of 10%.

Recent catches of Redbait East have been 4% of the 2020 estimate spawning biomass and below the Tier 2 exploitation rate for this stock of 5% (Smith et al. 2015). The total catch in 2020/21 in the SPF was 59% of the available TAC. On the basis of the information provided here, Redbait East is classified as sustainable. Patterson et al. (2021) also classified the Redbait East stock as 'not overfished' and 'not subject to overfishing'.

## 4.4.2 Redbait West

The main fishery for Redbait West is the SPF. The total SPF catch in 2020/21 was 0 t, down from 9 t in 2019/20 and 1,140 t in 2016/17. In 2019/20, the modal length for Redbait West was 190 mm FL, down from 240 mm FL in 2016/17, but data was only collected from one catch sample in 2019/20. In 2015/16, the catch was bimodal, with modes at 130 and 220 mm FL. Catches have mainly contained fish below the mean size at 50% maturity of ~250 mm FL.

Recent annual catches of Redbait West have not exceeded 9 t. The spawning biomass for Redbait West in 2017 was estimated to be 66,767 t (CI= 33,574–140,126; Ward et al. 2019). Recent catches of Redbait West have been <1% of this estimate of spawning biomass and below the Tier 1 exploitation rate for this stock of 10% (Smith et al. 2015). There was no catch reported in 2020/21: 0% of the available TAC. On the basis of the information provided here, Redbait West is classified as sustainable. Patterson et al. (2021) also classified the Redbait West stock as 'not overfished' and 'not subject to overfishing'.

# **5** Australian Sardine (*Sardinops sagax*)

# 5.1 Introduction

#### 5.1.1 Background to Fishery

Sardines (*Sardinops* spp.) form the basis of some of the world's largest fisheries (Schwartzlose et al. 1999) and have been the focus of extensive research (e.g. Stratoudakis et al. 2006, Ward et al. 2021b). In Australia, Sardine (*Sardinops sagax*) support several commercial fisheries in waters from southern Queensland to Western Australia (Ward and Staunton-Smith 2002, Piddocke et al. 2021).

Exploitation of Sardine in Australia has occurred since the 1800s (Kailola et al. 1993), but combined national catches did not exceed 1,000 t until the 1970s. Several purse seine fisheries developed in south-western Western Australia, and the annual catch in the west reached ~8,000 t in 1990 (Kailola et al. 1993). In 1991, a Sardine fishery was established in South Australia to provide fodder for the tuna mariculture industry (Ward and Staunton-Smith 2002).

In 1995 and 1998, two mass mortality events caused by Pilchard herpesvirus (Whittington et al. 2008) reduced the adult biomass of Australian Sardine populations by ~70% (Ward et al. 2001b). Catches in Western Australia have remained low since the mortality events (<3,000 t since 1999). In 2014, ~1,500 t of Australian Sardine were taken off Western Australia (Fletcher and Santoro 2015). The South Australian fishery grew quickly after the mortality events; in 2020, the total catch by the South Australian fishery was 41,295t (Grammer et al. 2021).

Off eastern Australia, the annual catch of Sardine increased rapidly from historical averages of 30-40 t to almost 5,000 t in 2008/09, but declined since then as a result of a reduction in fishing effort (Ward et al. 2014a, Izzo et al. 2017).

#### 5.1.2 Taxonomy

Australian Sardine (*Sardinops sagax*, Jenyns 1842) belong to the order Clupeiformes, which contains about 400 species in seven families, including *Clupeidae* (sardines, shads, menhadens, herrings) and *Engraulidae* (anchovies) (Eschmeyer and Fricke 2016).

The genus *Sardinops* has historically included five species: *S. ocellatus* off southern Africa; *S. neopilchardus* off southern Australia and New Zealand; *S. sagax* off the west coast of South America; *S. caeruleus* off the west coast of North America; and *S. melanostictus* off the Japanese coast (Whitehead 1985). Parrish et al. (1989) proposed the genus *Sardinops* is mono-specific with no valid sub-species, and that the name *Sardinops sagax* (Jenyns 1842) has taxonomic priority. This finding was confirmed by Grant and Leslie (1996). Grant et al. (1998) suggested that cluster and parsimony analyses of haplotypic divergences supported the hypothesis that there were three lineages within the genus: southern Africa (*ocellatus*) and Australia (*neopilchardus*); Chile (*sagax*) and California (*caeruleus*); and Japan (*melanostictus*). Currently, the accepted taxonomic classification for Sardine in the Indo-Pacific region is as one species, *Sardinops sagax*, with three confirmed linages within the genus.

The common name for Sardine off Australia has also varied through the years. It has been referred to as either the Australian Sardine or Pilchard. Since May 2006, Australian Sardine and *Sardinops sagax* (Jenyns 1842) have been listed in the Standard Fish Names List for Australia. *Sardinops sagax* is also the name used in the Australian Faunal Directory, Eschmeyer's Catalogue of Fishes of Australia, and FishBase. In this report, we use Sardine to refer to *S. sagax* in Australia and elsewhere.

## 5.1.3 Distribution

Sardine are found in waters off Australia, Japan, North and South America, Africa and New Zealand. In Australia, they occur throughout temperate waters between Rockhampton (Queensland) and Shark Bay (Western Australia), including northern Tasmania (Gomon et al. 2008).

## 5.1.4 Stock Structure

The Australian Sardine population has a complex stock structure. It is a meta-population with extensive mixing among adjacent sub-groups, illustrated by the rapid spread of Pilchard herpesvirus through the population in the late 1990s (Ward et al. 2001b, Whittington et al. 2008, Izzo et al. 2017). An integrated analysis of genetic, morphological, otolith, growth, reproductive and fishery data by Izzo et al. (2017) suggested the existence of least four stocks: 1) south-western Australia (Western Australia); 2) Great Australian Bight and Spencer Gulf (South Australia); 3) Victoria, Tasmania and southern NSW); and 4) eastern Australia (southern Queensland and northern NSW). The stock off south-western Australia appears to comprise two sub-stocks: west and south.

## 5.1.5 Movement

Sardines undergo extensive migrations. For example, schools of Sardine migrate north into waters off southern Queensland during winter-spring to spawn (Ward and Staunton-Smith 2002). Similarly, off Africa, Sardine migrate north and south along the coast to access conditions that are favourable for spawning and the survival of recruits (van der Lingen and Huggett 2003). The movement patterns of Sardines in Australian waters are poorly understood, although there is evidence of an ontogenetic shift in distribution in South Australia with larger, older fish most commonly found in shelf waters, and small, younger fish mainly found in embayments, including Spencer Gulf (Rogers and Ward 2007).

#### 5.1.6 Food and Feeding

Sardines have two feeding modes: filter-feeding on micro-zooplankton and phytoplankton, and particulate-feeding on macro-zooplankton. Sardines switch between the two modes depending on relative prey density (van der Lingen 1994, Louw et al. 1998). In South Australian waters, Sardines consume at least 12 different prey taxa; krill (29.6% biomass) and other unidentified crustaceans (22.2% biomass) are the major prey items (Daly 2007). Krill were found in greater numbers (65.3%) than other crustaceans (27%). Crab zoea, other decapods, copepods, polychaetes, fish eggs and larvae, and gelatinous zooplankton were also dietary components (Daly 2007).

#### 5.1.7 Age, Growth and Size

Sardines have been aged using growth increments in scales (Blackburn 1950) and otoliths (Butler et al. 1996, Fletcher and Blight 1996) and by modelling marginal increment formation in otoliths (Kerstan 2000). Several methods show that translucent zones form annually in otoliths of 1 year old fish off South Africa (Waldron 1998), 2 year olds off North America (Barnes et al. 1992) and 4 year olds off Western Australia (Fletcher and Blight 1996). Ageing sardine from southern Australia has been challenging due to difficulties associated with interpreting and counting growth zones (Rogers and Ward 2007).

Growth rates and maximum size of Australian Sardine vary in accordance with localised variation of food resources and environmental conditions (Ward and Staunton-Smith 2002). In southern Australia, Sardine rarely exceed 250 mm FL after 6 to 8 years (Rogers and Ward 2007). Larval and juvenile Sardine in southern Australian waters have growth rates of approximately 1.2 and 0.4 mm.day<sup>-1</sup>, respectively (Rogers and Ward 2007). Growth rates of Sardines were higher in South Australian waters than along other parts of the Australian coastline and lower than those in more productive boundary current ecosystems, such as the Benguela, Agulhas and Californian systems (Rogers and Ward 2007). A notable finding of the study was that fish in commercial catches were younger (and smaller) than those from fishery-independent samples.

#### 5.1.8 Reproduction

In Australia, Sardine usually spawn in open waters between the coast and shelf break (Blackburn 1950, Fletcher and Tregonning 1992, Fletcher et al. 1994). They are serial (batch) spawners with asynchronous oocyte development and indeterminate fecundity. The number of eggs released in a batch (batch fecundity) is correlated with female size and varies among locations and years (Lasker 1985). In South Australia, females spawn batches of ~18,000 pelagic eggs once about every 9 days during the extended spawning season (Ward et al. 2021b). In most locations, there is one spawning season per year, but off Albany in Western Australia, there are two (Fletcher 1990).

The peak spawning season is variable across the Australian distribution of Sardine. For example, in South Australia, spawning occurs during summer-autumn from January to

April (Ward et al. 2001c, Ward and Staunton-Smith 2002). Similarly, along the south coast of Western Australia spawning peaks between January and June (Gaughan et al. 2002), whereas Sardine off Fremantle have maximum GSI values during June (Murhling et al. 2008). Off southern Queensland and northern NSW, Sardine GSI values peak in winter to early spring (Ward and Staunton-Smith 2002, Ward et al. 2015c), whereas off southern NSW the peak occurs between July and December (Stewart et al. 2010, Ward et al. 2011a). Off Victoria, GSI values of Sardine are greatest from spring to early summer (Hoedt and Dimmlich 1995, Neira et al. 1999).

The size and age at of sexual maturity in Sardine varies between locations, and ranges from 100 to 180 mm FL and 1.8 to 2.8 years (Blackburn 1950, Fletcher 1990, Staunton-Smith and Ward 2000). In South Australia, 50% of males are sexually mature at 139 mm FL and females at 144 mm FL (Ward et al. 2020c).

## 5.1.9 Early Life History and Recruitment

Sardine have a relatively long larval phase: eggs hatch approximately two days after fertilization and yolk-sac larvae are 2.2–2.5 mm TL (Neira et al. 1998). Larvae metamorphose at 1–2 months of age and at lengths of 35–40 mm TL. Larvae are known to undertake vertical migrations to prevent passive transport away from regions with favourable environmental conditions for survival (Watanabe et al. 1996, Logerwell et al. 2001, Curtis 2004). Survival rates of sardine eggs and larvae strongly affect recruitment success (Louw et al. 1998). Large variations in abundance that characterise sardine populations worldwide are attributed to fluctuations in recruitment, which can be influenced by environmental factors, regime shifts and over-fishing (Galindo-Cortes et al. 2010). Larval survival is a key determinant of recruitment success, but factors affecting survivorship may vary spatially and temporally. The effects of food availability on larval survival have been discussed at length (Galindo-Cortes et al. 2010), but there has been less consideration about how predation on eggs and larvae may affect recruitment success.

In South Australia, Sardine larvae are highly abundant at temperature and salinity fronts that form near the mouths of the two Gulfs during summer and autumn (Bruce and Short 1990) and in mid-shelf waters of the eastern and central Great Australian Bight (Ward et al. 2020c, Grammer et al. 2021). Juvenile Sardine occupy nursery areas that include shallow embayments and semi-protected waters. The factors affecting recruitment success of Sardine are poorly understood.

## 5.1.10 Stock Assessment

The DEPM was developed to assess the status of northern anchovy (*Engraulis mordax*) stocks off the coast of California (Parker 1980, Lasker 1985) and is widely used for assessing spawning-stock biomass of sardine worldwide (see review in Barangé et al. 2009), e.g. Atlanto-Iberian Sardine (*Sardina pilchardus*) (Bernal et al. 2011a, Bernal et al. 2011b). This approach provides direct estimates of spawning biomass for the basis of

management decisions. The DEPM has been used extensively to estimate the spawning biomass of Australian Sardine in South Australia since 1995 (Ward et al. 2021b).

The DEPM has been applied to the eastern stock of Sardine off east Australia in 2014 (Ward et al. 2015b) and 2019 (Ward et al. 2021a). Prior to 2014, a preliminary DEPM was applied to the eastern stock of Sardine in 2004 using samples collected along the southern Queensland and northern NSW coast during a survey of Blue Mackerel (Ward et al. 2007). The DEPM has been applied to the eastern component of the south-eastern Sardine stock in 2014 (Ward et al. 2015a) and 2019 (Ward et al. 2022) and to the western component in 2016/17 (Ward et al. 2018).

## 5.1.11 Recreational fishing

Information on the magnitude of recreational catches of Sardine is not available. The most recent report on the Status of Australian Fish Stocks indicated that recreational and indigenous catches of Australian Sardine are likely to be negligible (Piddocke et al 2021).

## 5.1.12 Biomass Estimates

#### East

The DEPM was applied to the eastern stock of Australian Sardine East in 2004 (Ward et al. 2009), 2014 (Ward et al. 2015b) and 2019 (Ward et al. 2021a). The preliminary estimate of spawning biomass for Australian Sardine East in 2004, calculated from the 'best' estimate of each parameter, was 28,809 t (Ward et al. 2007). 'Minimum' and 'maximum' estimates ranged from 9,161 to 58,673 t. The 'best' estimate was considered conservative and likely negatively biased as spawning season and area varies temporally and spatially on the east coast of Australia (Ward and Staunton-Smith 2002). It is unlikely that the entire spawning area was sampled during peak spawning season (Ward et al. 2007).

The first dedicated DEPM survey for Australian Sardine East was undertaken in August/September 2014 off eastern Australia; the estimate of spawning biomass for the survey was 49,575 t (95% CI = 24,179 - 213,323) (Ward et al. 2015b). The estimated spawning area off eastern Australia was 22,400 km<sup>2</sup>, comprising 34.2% of the total area sampled (65,528 km<sup>2</sup>) (Ward et al. 2015b). Live Sardine eggs (n = 3,461) were collected from 89 of the 262 (34.0 %) stations between Sandy Cape, Queensland to just south of Newcastle, NSW. Mean daily egg production ( $P_0$ ) was 52.6 eggs·day<sup>-1</sup>·m<sup>-2</sup>. The highest densities of Sardine eggs were recorded in waters off southern Queensland from Fraser Island to Caloundra, between Byron Bay and Coffs Harbour in northern NSW and off Port Stephens where SSTs ranged between 18 and 21°C.

The second dedicated DEPM survey for Australian Sardine East was undertaken in September 2019 off eastern Australia; the estimate of spawning biomass for the survey was 42,724 t (95% CI = 15,487–69,962) (Ward et al. 2021a). The estimate of spawning area off eastern Australia was 14,281 km<sup>2</sup>, which was 22.9% of the total area sampled

(62,476 km<sup>2</sup>) (Ward et al. 2021a). Live Sardine eggs (n = 4,667) were collected at 58 of the 251 sites (23.1%) between southern Queensland and central NSW.  $P_0$  was 53.9 eggs.day<sup>-1</sup>.m<sup>-2</sup>. High egg densities were mostly found on the mid to outer shelf off southern Queensland, at inshore to mid shelf sites in northern NSW, and at mainly inshore sites off central NSW; SSTs ranged between 16 and 22°C.

The estimates of spawning biomass for the eastern stock of Australian Sardine East for 2014 and 2019 are considered to be reliable, as they are based on robust and/or conservative estimates of all key parameters (Ward et al. 2015b, 2021a). The 14% decline in the spawning biomass of Sardine from 49,575 t in 2014 to 42,724 t in 2019, combined with the 36% decline in spawning area over the same period, suggest that stock abundance has declined over this period (Ward et al 2021a). Because of the low level of recent catches in this sub-area (<800 t; Figure 5-1), it seems likely that this decline reflects a natural fluctuation in abundance rather than a fishing induced impact (Ward et al. 2021a).

## 5.1.13 Management Strategy Evaluation

Smith et al. (2015) concluded the harvest rate of 15% may be too low for Australian Sardine East and suggested a Tier 1 harvest rate of 33%, with the Tier 1 rate being applied for not more than 5 years. Tier 2 harvest rates for Australian Sardine East were recommended to be 50% of Tier 1 rates and should not be applied for more than 5 years. The study results also indicated it is not safe to apply Tier 2 harvest rates unchecked for long periods of time (Smith et al. 2015). The Atlantis-SPF biomass estimate for Australian Sardine East is 147,000 t (typical range: 32,000–184,000 t) (Smith et al. 2015).

## 5.1.14 Management

Sardine in the Australian Sardine sub-area of the East sub-area in the SPF are currently managed at the Tier 1 under the Harvest Strategy. DEPM assessments of the eastern stock of Australian Sardine have been conducted for the East sub-area of the SPF in 2004, 2014 and 2019

## 5.2 Methods

## 5.2.1 Fishery Statistics

Fishery statistics from 1984/85 to 2020/21 have been supplied by relevant jurisdictions and collated by SARDI Aquatic Sciences. Annual data are reported in fishing seasons (May 1 to April 30) rather than financial years as was done in previous assessments (e.g. Ward et al. 2013, 2014c, 2015c). Data reported for Australian Sardine East include the Australian Sardine sub-area and not the entire East sub-area, due to the eastern and south-eastern Sardine stock separations that occur in southern NSW/eastern VIC (Figure 1-1; see Izzo et al. 2017).

Estimates of total annual catch supplied for Australian Sardine East include data from the NSW Ocean Fisheries (Hauling, Trap and Line, Trawl), NSW Estuary Fisheries (General and Prawn Trawl) and the Commonwealth SPF.

Mean annual CPUE of Australian Sardine East in the Commonwealth SPF is calculated for purse-seine (tonnes net-set<sup>-1</sup> ±SE) from 2000/01 to 2020/21.

#### 5.2.2 Biological Information

Length-frequency data for Australian Sardine sampled from commercial catches taken in NSW were supplied by NSW DPI from 2004/05 to 2016/17. Additional samples were collected from commercial catches taken from the north (Iluka) and south-central (Eden) coast of NSW between March 2009 and January 2010 for biological analysis. These fish were dissected and morphometric data collected by NSW DPI, while otoliths were interpreted for age by SARDI Aquatic Sciences using the methods of Rogers and Ward (2007).

AFMA observers collected biological samples of Sardine during trips in September 2012 and August 2013 which were supplied to SARDI Aquatic Sciences for processing. The fish were measured (mm FL), and a random sub-sample was retained for ageing. Australian Sardines were also sampled from commercial catches taken in NSW between July 2013 and August 2014 to determine population size and age structures and monitor reproductive activity.

Ages were derived from the otolith weights in 2012/13 using the relationship calculated from South Australian commercial catches (Rogers and Ward 2007). In all other years, ages were based annual growth increment counts in whole otoliths.

Summarised biological data from NSW DPI are presented in financial years and all other biological data are presented in fishing seasons (1 May to 30 April).

The number and spatio-temporal coverage of the biological samples of Australian Sardine East are limited in comparison to the magnitude of catches since the mid-1990s, and may not provide a good representation of the catch.

## 5.3 Results

#### 5.3.1 Australian Sardine East

## 5.3.1.1 Fishery Statistics

#### Number of vessels

Prior to 1999/00, >85 vessels annually reported catches of Australian Sardine East; 99% were from NSW. Since then, vessel numbers taking Australian Sardine East declined to 15 vessels in 2012/13 and have averaged 27 per year from 2013/14 to 2018/19 (80% from

#### Securing Australia's fishing future

NSW). Number of vessels declined to 19 in 2019/20 and 15 in 2020/21 (70% from NSW and 24% from SPF).

#### Annual patterns: Total catch

Total catches of Australian Sardine have varied over the last 34 years (Figure 5-1a). Catches were low in the mid- to late-1980s (<200 t), increased to ~450 t in 1996/97 and fell to low levels in the early 2000s. The low catches of the late 1990s and early 2000s resulted from the widespread die-off of Sardine from the Pilchard herpesvirus that spread through the Australian population in 1995 and 1998 (Ward et al. 2001b, Whittington et al. 2008). The stocks recovered and catches increased to 3,761 t in 2007/08 (Figure 5-1a). Total catches declined to 238 t in 2012/13 and have remained below 650 t until 2018/19. Total catch was 733 t in 2019/20 and 717 t in 2020/21. The main fisheries taking Australian Sardine in the Sardine sub-area are the NSW Ocean Hauling Fishery (88% of 2020/21 catch) and the SPF.

#### Annual patterns: Catch, Effort and CPUE

Purse seining has been the main method used to take Australian Sardine East. In the SPF, catches of Australian Sardine East have been solely taken by purse seine (Figure 5-1b, c). Historically, annual purse seining effort in the SPF has been substantially less than in the NSW fishery but with only slightly lower catch. For example, in 2007/08, effort and catch in the NSW fishery was 1,948 t with 354 boat days compared to 1,759 t and 56 netsets (49 boat days) in the SPF (Figure 5-1b, d). Catch trends in the SPF are similar to those in the NSW fishery, with the highest catches in the mid to late 2000s (Figure 5-1 b, d). Temporal effort in the SPF was relatively stable, averaging 53 net-sets annually from 2005/06 to 2011/12. Both effort and catch of Australian Sardine East in the SPF decreased to low levels between 2011/12 and 2013/14 (Figure 5-1b). Fishing effort and catch increased in 2014/15 (57 net-sets, catch: 152 t) and remained steady through 2016/17 (effort: 57 net-sets, catch: 131 t). Effort was 61 net-sets in 2017/18 (99 t of catch), 81 net-sets in 2018/19 (132 t of catch) and 90 net-sets in 2019/20 (223 t of catch). Effort decreased to 32 net-sets (82 t of catch) in 2020/21 (Figure 5-1b).

Mean annual CPUE of purse seining in the SPF for Sardine declined from 31 t·net-set<sup>-1</sup> in 2007/08 to 7 t·net-set<sup>-1</sup> in 2010/11 (Figure 5-1c). Mean annual CPUE has been <3 t·net-set<sup>-1</sup> from 2011/12 to the present.

In the NSW fishery, purse seining effort for Australian Sardine has been highly variable among years; in 2000/01, effort was 108 boat days with a catch of only 18 t (Figure 5-1d). Fishing effort remained low in the early 2000s and increased to a peak of >350 boat days in the late 2000s. Catches also peaked at ~1,950 t during that time (Figure 5-1d). From 2014/15 to 2016/17, effort averaged 121 boat days and 429 t of catch per year. In 2018/19, effort was 137 boat days with 457 t of catch. Effort increased in 2019/20 to 203 boat days with 490 t of catch. Annual catch and effort data are confidential in the NSW fishery from 2011/12–2013/14, 2017/18 and 2022/21 due to the low number (<6) of licence holders reporting catch.



Figure 5-1. Fishery statistics for Australian Sardine East (Sardine Sub-area only). (A) Total annual catch (tonnes) for all jurisdictions by fishing season from 1984/85 to 2020/21. Long-term purse seining trends in the SPF from 2000/01 to 2020/21: (B) annual landed catch (tonnes) and effort (net-sets); (C) mean annual CPUE (t-net-set<sup>-1</sup>;  $\pm$ SE). Long term purse seining trends in the NSW Ocean Hauling Fishery: (D) annual landed catch (tonnes) and effort (tonnes) and effort (boat days). (**x**) indicates data confidentiality where <6 licence holders reported landings.

# 5.3.1.2 Biological Information

Length-frequency data were collected from 18,610 Australian Sardine East sampled from purse seine catches in the NSW Ocean Hauling Fishery between 2004/05 and 2016/17 (Table 5-1). The number of samples collected in each year ranged between 2 (2004/05) and 54 (2009/10).

In the SPF, length-frequency data were collected from 4,068 Australian Sardine East taken by purse seining off NSW between 2009/10 and 2014/15 (Table 5-2). Of these samples, age-frequency data were collected from 1,215 individuals (Table 5-2).

Table 5-1. Summary of Australian Sardine East samples collected from commercial NSW State purse seine catches between 2004/05 and 2016/17 (data supplied by NSW DPI).

Season	No. of samples	No. of fish	Size range (mm FL)	
2004/05	2	249	90–210	
2005/06	7	592	80–240	
2006/07	31	3,098	70–230	
2007/08	12	1,209	90–230	
2008/09	8	860	110–210	
2009/10	54	5,579	50–230	
2010/11	5	473	100–220	
2011/12	6	691	100–200	
2012/13	4	538	100–180	
2013/14	30	2,075	120–190	
2014/15	12	1,223	120–200	
2015/16	12	1,186	90–200	
2016/17	8	733	120–210	

Table 5-2. Summary of Australian Sardine East catch samples collected from commercial SPF landings off NSW by purse seine from 2009/10 to 2014/15.

Season	NSW Region	No. of samples	Length- frequency n	Age- frequency n	Size range (mm FL)	Age range (years)
2009/10	North	15	240	155	120–190	0–3
2009/10	South	6	330	167	127–213	0–5
2012/13	North	3	208	32	120–175	1–3
2013/14	North	40	1,840	492	68–175	0—5
2014/15	North	32	1,450	369	124–195	0—6

#### Size Structure

Annual size distributions for Australian Sardines sampled between 2004/05 and 2016/17 from the NSW Ocean Hauling Fishery were mainly between 100 and 200 mm FL, although the modal length varied among years (Figure 5-2). For some years (i.e. 2004/05 to 2006/07, 2011/12), the size structure was bimodal with a dominant mode at ~130 to 140 mm FL (i.e. the approximate size at sexual maturity for Australian Sardine) and a smaller mode at ~170 to 190 mm FL. In all other years, the size structures contained a single dominant mode between 130 and 180 mm FL. The length mode has increased since 2012/13 from 140–150 mm FL to 180 mm FL in 2016/17 (Figure 5-2).



Figure 5-2. Length-frequency distributions of Australian Sardine East sampled from purse seine catches taken in NSW from 2004/05 to 2016/17. Data supplied by NSW DPI; n = number of fish. See Table 5-1 for sample N.

On average, the size structures of catches off NSW in 2009/10 of Australian Sardine East were larger in the south than in the north (Figure 5-3). Purse seine catch samples taken in the SPF between 2012/13 and 2014/15 indicated an increase in the size structure over time but a consistent modal length at 150 mm FL (Figure 5-3).



Figure 5-3. Length-frequency distributions of Australian Sardine East caught in the SPF by purse seine along the northern and southern NSW coastline; n = number of fish. See Table 5-2 for sample N.

#### Age structure

Ages of Australian Sardine East collected from purse seine catches in the SPF along the south coast of NSW in 2009/10 ranged from 0+ to 5 years, whereas catches from the northern region contained fish aged 0+ to 3 years (Figure 5-4). Commercial catch samples from the north in 2012/13 mostly consisted of fish aged 2 years. In 2013/14 and 2014/15, the range of ages in the northern region continued to increase, with more fish in the 3 and 4 year old age classes. In 2014/15, the maximum estimated age of fish was 6 years (Figure 5-4).



Figure 5-4. Age-frequency distributions of Australian Sardine East caught by purse seine in the SPF along the northern and southern NSW coastline; n = number of fish. See Table 5-2 for sample N.

## 5.4 Summary and Conclusions

#### 5.4.1 Australian Sardine East

The main fisheries for Australian Sardine in the Sardine sub-area are the NSW Ocean Hauling Fishery and the SPF. Total annual catches peaked at 3,761 t in 2007/08, declined to 239 t in 2012/13, increased to 733 t in 2019/20, and was 717 t in 2020/21. Catches were mainly taken by purse seine, and since 2000/01, annual fishing effort in the SPF has been substantially less than in the NSW Ocean Hauling Fishery, but with only slightly lower catch. Both effort and catch of Sardine in the SPF decreased to low levels between 2011/12 and 2013/14, increased through 2019/20, and have again declined in 2020/21.

There has been a general reduction in total catch since 2008/09 that reflects a significant reduction in the size of the fishing fleet. Less than 40 vessels annually have reported catch since 2011/12, compared to >90 vessels prior to 1999. Other factors that may have contributed to the reduction in catch and effort include a fire in a major fish processing factory in Eden (southern NSW) and apparent movement of Sardines from inshore to offshore waters (AFMA 2014).

Size structures of Australian Sardine East in purse seine catches since 2004/05 have varied among years, with the modal length ranging from 130–180 mm FL. The length mode of purse seine catches increased from 150 mm FL in 2012/13 to 180 mm FL in 2016/17. Sardine in catches have been at or above the mean size at 50% maturity of ~150 mm FL (Ward and Rogers 2007) with age classes of mainly 2-4 years. Age and length structures are difficult to interpret due to changes in fishing effort over time, differing ageing methods and regional stock differences (e.g. northern vs. southern NSW).

The spawning biomass of the eastern stock of Australian Sardine East was estimated to be 49,575 t in 2014 (Ward et al. 2015b) and 42,724 t in 2019 (Ward et al. 2021a). The spawning area of the eastern stock reduced by 36% between the 2014 and 2019 (14% change in spawning biomass). Spawning area is strongly correlated with adult abundance in many species of small pelagic fishes including Sardine (e.g. Barangé, et al. 2009; Ward et al. 2021b), indicating the population may have declined over that period (Ward et al. 2021a). It is unlikely that this apparent decline in abundance has been driven primarily by fishing pressure, because total catches of Sardine have been <800 t since 2010/11 and <2 % of the spawning biomass in 2019 (Ward et al. 2021a). This apparent change in abundance may be due to natural fluctuations in recruitment and mortality which are known to occur in this species (e.g. Barangé et al. 2009; Ward et al. 2020c).

Current catches of Australian Sardine East in the Sardine sub-area are assessed as sustainable, as they are <2% of the estimated spawning biomass for 2014 (Ward et al. 2015b), and below the Tier 1 exploitation rate for this stock of 20% (Smith et al. 2015). Piddocke et al. (2021) classified the biological stock of Australian Sardine East as sustainable. The Australian Sardine stock in the East is classified as 'not overfished' and 'not subject to overfishing' by Patterson et al. (2021).

# 6 General Summary and Conclusions

The most recent classifications of the stock status of Jack Mackerel East and West, Blue Mackerel East and West, and Australian Sardine East concluded that the biological stocks are sustainable (Piddocke et al. 2021). Jack Mackerel, Blue Mackerel and Redbait in both sub-areas, and Australian Sardine in the Sardine Sub-area were classified as 'not over-fished' and 'not subject to overfishing' by Patterson et al. (2021). The evidence presented in this report confirms that recent catches of Blue Mackerel, Jack Mackerel and Redbait in the East and West sub-areas and Australian Sardine in the Sardine in the Sardine sub-area have been sustainable.

DEPM surveys have now been conducted for all SPF stocks. All stocks in the East subarea have undergone a DEPM survey in the last 5 years and are at Tier 1 of the Harvest Strategy. In the West sub-area, DEPM surveys for Jack Mackerel and Rebait have occurred in the last 5 years and these stocks are also at Tier 1. The last DEPM survey for Blue Mackerel West was in 2005 and this stock is at Tier 3. Surveys conducted in the West sub-area have only covered a limited portion of that region. As a result, the distribution and abundance of SPF species in the West sub-area in the region west of Kangaroo Island is poorly understood. The estimates of spawning biomass now available could be used to inform the establishment of preliminary target (e.g. B<sub>50</sub>) and limit reference points (e.g. B<sub>20</sub>) for each stock in the SPF. These reference points could be revised as new information becomes available.

Mounting evidence, resulting from spawning biomass surveys, is negating the idea of an East-West stock structure for Jack Mackerel and Redbait, where Bass Strait is the natural barrier between stocks due to the prevailing ocean currents (e.g. Bulman 2008). Egg distributions from spawning biomass surveys for Jack Mackerel in 2014, 2016/17 and 2019 suggest a break in stock structure may occur along the Bonney Coast of South Australia (Ward et al. 2015a, 2018, 2020a). The main spawning stock of Jack Mackerel for south-eastern Australia is thought to be located from western Victoria through Bass Strait to eastern Tasmania and southern NSW (Bulman et al. 2015, Ward et al. 2018, 2020a). Egg distributions from spawning biomass surveys for Redbait in 2005, 2006, 2017 and 2020 suggest Redbait in south-eastern Australian may be a continuous stock along the continental shelf break and inner slope from southern NSW, around Tasmania to Kangaroo Island in South Australia (Neira et al. 2008, Ward et al. 2019, Grammer et al. 2022). There was no evidence of spawning activity of Redbait through Bass Strait (Ward et al. 2019, Grammer et al. 2022).

Fishery-dependent data have been assessed to evaluate the potential for changes in stock status between DEPM surveys. The spatial and temporal distribution of catches have varied over time, driven mainly by changes in the fishing operations. Differences observed in age and length structures among years and sub-areas, have been difficult to interpret due to these changes in the fishery. Since 2014/15, changes in CPUE and size and age structures may have been driven more by differences in vessels and their operations in the SPF, rather than from fishing pressure.

Securing Australia's fishing future

# References

- AFMA. 2008. Small Pelagic Fishery Harvest Strategy (last revised April 2017). Australian Fisheries Management Authority, Canberra. 9 pp.
- AFMA. 2009. Small Pelagic Fishery Management Plan 2009. Australian Fisheries Management Authority, Canberra, Federal Register of Legislative Instruments F2010L00081. 51 pp.
- AFMA. 2014. Small Pelagic Fishery Management Arrangements Booklet 2014, Australian Fisheries Management Authority. Canberra, Australia. 32 pp.
- Arneri, E., P. Carpi, F. Donatato, and A. Santojanni. 2011. Growth in small pelagic fishes and its implications in their population dynamics. Biologia Marina Mediterranea **18**: 106-113
- Barangé, M., M. Bernal, M. C. Cercole, L. Cubillos, C. L. Cunningham, G. M. Daskalov, J. A. A. De Oliveira, and e. al. 2009. Current trends in the assessment and management of small pelagic fish stocks. *in* D. M. Checkley, C. Roy, J. Alheit, and Y. Oozek, editors. Climate Change and Small Pelagic Fish. Cambridge University Press, Cambridge, UK.
- Barnes, J. T., A. D. Maccall, L. D. Jacobson, and P. Wolf. 1992. Recent Population Trends and Abundance Estimates for the Pacific Sardine (*Sardinops-Sagax*). California Cooperative Oceanic Fisheries Investigations Reports **33**:60-75.
- Bernal, M., Y. Stratoudakis, S. Wood, L. Ibaibarriaga, A. Uriarte, L. Valdés, and D. Borchers. 2011a. A revision of daily egg production estimation methods, with application to Atlanto-Iberian sardine. 1. Daily spawning synchronicity and estimates of egg mortality. ICES Journal of Marine Science: Journal du Conseil 68:519-527.
- Bernal, M., Y. Stratoudakis, S. Wood, L. Ibaibarriaga, L. Valdés, and D. Borchers. 2011b. A revision of daily egg production estimation methods, with application to Atlanto-Iberian sardine. 2. Spatially and environmentally explicit estimates of egg production. ICES Journal of Marine Science: Journal du Conseil 68:528-536.
- Blaber, S. J. M., and C. M. Bulman. 1987. Diets of fishes of the upper continental slope of eastern Tasmania: content, calorific values, dietary overlap and trophic relationships. Marine Biology 95:345-356.
- Blackburn, M. 1950. Studies on the age, growth, and life history of the sardine *Sardinops neopilchardus* (Steindachner), in Southern and Western Australia. Australian Journal of Marine and Freshwater Research **1**:221-258.
- Browne, A. 2005. Changes in the growth and age structure of jack mackerel in south-east Australian waters. Honours Thesis, University of Tasmania, School of Zoology. 58 pp.
- Bruce, B. D., and D. A. Short. 1990. Observations on the distribution of larval fish in relation to a frontal system at the mouth of Spencer Gulf, South Australia. Bureau of Rural Resources Proceedings **15**:124-137.
- Bulman, C., F. Althaus, X. He, N. J. Bax, and A. Williams. 2001. Diets and trophic guilds of demersal fishes of the south-eastern Australian shelf. Marine and Freshwater Research **52**:537-548.
- Bulman, C., S. Condie, J. Findlay, B. Ward, and J. Young. 2008. Management zones from small pelagic fish species stock structure in southern Australian waters. Final Report for FRDC Project 2006/076. CSIRO Marine and Atmospheric Research: Hobart, Tasmania.
- Bulman, C., S. Davenport, and F. Althaus. 2000. Trophodynamics. In: Habitat and fisheries production in the South East Fishery ecosystem. Final report to the Fisheries Research and Development Corporation, Project No. 94/040, (eds N.J. Bax and A. Williams). Division of Marine Research, CSIRO Marine Laboratories, Hobart.
- Bulman, C. M., S. A. Condie, F. J. Neira, S. D. Goldsworthy, and E. A. Fulton. 2011. The trophodynamics of small pelagic fishes in the southern Australian ecosystem and the

implications for ecosystem modelling of southern temperate fisheries. Final report for FRDC project 2008/023. CSIRO Marine and Atmospheric Research. Hobart. 101 pp.

- Bulman, C. M., E. A. Fulton, and A. D. M. Smith. 2015. Jack mackerel stock structure in the SPF. CSIRO Marine and Atmospheric Research. Hobart, Tasmania. 19 pp.
- Butler, J. L., M. L. Granados, J. T. Barnes, M. Yaremko, and B. J. Macewicz. 1996. Age composition, growth, and maturation of the Pacific sardine (*Sardinops sagax*) during 1994. California Cooperative Oceanic Fisheries Investigations Reports **37**:152-159.
- Catanese, G., M. Manchado, and C. Infante. 2010. Evolutionary relatedness of mackerels of the genus Scomber based on complete mitochondrial genomes: Strong support to the recognition of Atlantic *Scomber colias* and Pacific *Scomber japonicus* as distinct species. Gene **452**:35-43.
- Choat, J. H., and L. M. Axe. 1996. Growth and longevity in acanthurid fishes; an analysis of otolith increments. Marine Ecology Progress Series **134**:15-26.
- Curtis, K. A. 2004. Fine scale spatial pattern of Pacific sardine (*Sardinops sagax*) and northern anchovy (*Engraulis mordax*) eggs. Fisheries Oceanography **13**:239-254.
- Daly, K. 2007. The diet and guild structure of the small pelagic fish community in the eastern Great Australian Bight, South Australia. Honours thesis, University of Adelaide.
- Dennis, J., G. Grammer, T. Ward, J. Smart, and C. Huveneers. 2021. Using otolith chronologies to identify extrinsic drivers of growth of two commercially-targeted small pelagic fish species. Fishery Bulletin **119**:135-148.
- Emery, T., J. Bell, J. Lyle, and K. Hartmann. 2015. Tasmanian Scalefish Fishery Assessment 2013/14. Institute for Marine and Antarctic Studies, University of Tasmania. 102pp.
- Eschmeyer, W. N., and R. Fricke. 2016. Catalog of Fishes: Genera, Species, References. (http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp). Electronic version accessed 28 January 2016.
- Ewing, G. P., and J. M. Lyle. 2009. Reproductive dynamics of redbait, Emmelichthys nitidus (Emmelichthyidae), from south-eastern Australia. Fisheries Research **97**:206-215.
- Ewing, G. P., J. M. Lyle, R. J. Murphy, J. M. Kalish, and P. E. Ziegler. 2007. Validation of age and growth in a long-lived temperate reef fish using otolith structure, oxytetracycline and bomb radiocarbon methods. Marine and Freshwater Research **58**:944-955.
- Fisheries New Zealand. 2021. Fisheries Assessment Plenary, May 2021: stock assessments and stock status. Compiled by the Fisheries Science Team, Fisheries New Zealand, Wellington, New Zealand. 1782 p.
- Fletcher, W., and R. Tregonning. 1992. Distribution and timing of spawning by the Australian pilchard (*Sardinops sagax neopilchardus*) off Albany, Western Australia. Marine and Freshwater Research **43**:1437-1449.
- Fletcher, W. J. 1990. A synopsis of the biology and exploitation of the Australian sardine, Sardinops neoplichardus (Steindachner). I: Biology. Western Australian Department of Fisheries. Fisheries Research Report.
- Fletcher, W. J., and S. J. Blight. 1996. Validity of using translucent zones of otoliths to age the pilchard *Sardinops sagax neopilchardus* from Albany, western Australia. Marine and Freshwater Research **47**:617-624.
- Fletcher, W. J., and K. Santoro. 2015. Status Reports of the Fisheries and Aquatic Resources of Western Australia 2014/15: The State of the Fisheries. Department of Fisheries, Western Australia. 353 pp.
- Fletcher, W. J., R. J. Tregonning, and G. J. Sant. 1994. Interseasonal variation in the transport of pilchard eggs and larvae off southern Western Australia. Marine Ecology Progress Series 111:209-224.

- Fowler, A. J., and D. Short. 1998. Validation of age determination from otoliths of the King George whiting *Silliginodes punctata* (Perciformes). Marine Biology **130**:577-587.
- Fulton, E. A. 2013. Simulation analysis of jack mackerel stock size: Ecosystem model based plausibility study. CSIRO, Australia. 24 pp.
- Furlani, D., A. Williams, and N. Bax. 2000. Fish biology (length and age). In: Habitat and fisheries production in the South East Fishery ecosystem. Final report to the Fisheries Research and Development Corporation, Project No. 94/040, (eds N.J. Bax and A. Williams). Division of Marine Research, CSIRO Marine Laboratories, Hobart.
- Galindo-Cortes, G., J. A. De Anda-Montañez, F. Arreguín-Sánchez, S. Salas, and E. F. Balart. 2010. How do environmental factors affect the stock–recruitment relationship? The case of the Pacific sardine (*Sardinops sagax*) of the northeastern Pacific Ocean. Fisheries Research 102:173-183.
- Garrido, S., A. Marçalo, J. Zwolinski, and C. D. van der Lingen. 2007. Laboratory investigations on the effect of prey size and concentration on the feeding behaviour of *Sardina pilchardus*. Marine Ecology Progress Series **330**:189-199.
- Gaughan, D. J., W. J. Fletcher, and J. P. McKinlay. 2002. Functionally distinct adult assemblages within a single breeding stock of the sardine, *Sardinops sagax*: management units within a management unit. Fisheries Research **59**:217-231.
- Gaughan, D. J., and R. W. D. Mitchell. 2000. The biology and stock assessment of the tropical sardine, Sardinells lemura, off the mid-west coast of Western Australia., Fisheries Western Australia. No. 119, North Beach.
- Giannini, F., P. I. Hobsbawn, G. A. Begg, and M. Chambers. 2010. Management strategy evaluation (MSE) for the harvest strategy for the Small Pelagic Fishery. FRDC Project 2008/064, Final Report. Fisheries Research and Development Corporation and Bureau of Rural Sciences, Canberra. 157 pp.
- Giri, K., and K. Hall. 2015. South Australian Recreational Fishing Survey 2013/2014. Fisheries Victoria Internal Report Series No. 62.
- Gomon, M., D. Bray, and R. H. Kuiter. 2008. Fishes of the Australia's South Coast. Museum of Victoria. New Holland Publishers (Australia).
- Gonçalves, P., A. M. Costa, and A. G. Murta. 2009. Estimates of batch fecundity and spawning fraction for the southern stock of horse mackerel (*Trachurus trachurus*) in ICES Division IXa. ICES Journal of Marine Science: Journal du Conseil **66**:617-622.
- Grammer, G., F. Bailleul, A. Ivey, and J. Smart. 2021. Stock assessment of Australian Sardine (*Sardinops sagax*) off South Australia 2021. Report to PIRSA Fisheries and Aquaculture. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2007/000765-8. SARDI Research Report Series No. 1120. 96pp.
- Grammer, G. L., T. M. Ward, A. R. Ivey, and J. P. Keane. 2022. Spawning biomass of Redbait (*Emmelichthys nitidus*) in the East sub-area of the Small Pelagic Fishery during October 2020. Report to the Australian Fisheries Management Authority. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. .
- Grant, W. S., and B. W. Bowen. 1998. Shallow population histories in deep evolutionary lineages of marine fishes: Insights from sardines and anchovies and lessons for conservation. Journal of Heredity **89**:415-426.
- Grant, W. S., and R. W. Leslie. 1996. Late Pleistocene dispersal of Indian-Pacific sardine populations in an ancient lineage of the genus Sardinops. Marine Biology **120**:133-142.
- Harris, G. P., F. B. Griffiths, and L. A. Clementson. 1992. Climate and the fisheries off Tasmania interactions of physics, food chains and fish. South African Journal of Marine Science **12**:585-597.

- Heemstra, P. C., and J. E. Randall. 1977. A revision of the Emmelichthyidae (Pisces: Perciformes). Marine and Freshwater Research **28**:361-396.
- Henry, G. W., and J. M. Lyle. 2003. The National Recreational and Indigenous Fishing Survey. Final Report of FRDC Project No. 99/158. Final Report to FRDC 1999/158. New South Wales Fisheries Research Centre. 188 pp.
- Hoedt, F. E., and W. F. Dimmlich. 1995. Egg and Larval Abundance and Spawning Localities of the Anchovy (*Engraulis australis*) and Pilchard (*Sardinops neopilchardus*) near Phillip Island, Victoria. Marine and Freshwater Research **46**:735-743.
- Horn, P. L., H. L. Neil, L. J. Paul, and P. J. McMillan. 2012. Age verification, growth and life history of rubyfish *Plagiogeneion rubiginosum*. New Zealand Journal of Marine and Freshwater Research 46:353-368.
- Infante, C., E. Blanco, E. Zuasti, A. Crespo, and M. Manchado. 2006. Phylogenetic differentiation between Atlantic *Scomber colias* and Pacific *Scomber japonicus* based on nuclear DNA sequences. Genetica:1-8.
- Izzo, C., T. M. Ward, A. R. Ivey, I. M. Suthers, J. Stewart, S. C. Sexton, and B. M. Gillanders. 2017. Integrated approach to determining stock structure: implications for fisheries management of sardine, *Sardinops sagax*, in Australian waters. Reviews in Fish Biology and Fisheries 27:267-284.
- Jordan, A. R. 1994. Age, growth and back-calculated birthdate distributions of larval jack mackerel *Trachurus declivis* (Pisces: Carangidae), from eastern Tasmanian coastal waters. Journal of Marine and Freshwater Research 45:19-33.
- Lindholm, R., and J. G. H. Maxwell. 1988. Stock separation of jack mackerel Trachurus declivis (Jenyns, 1841), and yellowtail T. novaezealandiae (Richardson, 1843) in southern Australian waters using principal component analysis. CSIRO Marine Laboratories Report No. 189. Hobart, Tasmania. 7 pp.
- Shuntov, V. P. 1969. Some features of the ecology of pelagic fishes in the Great Australian Bight. Problems of Ichthyology 9:801-809.
- Smolenski, A. J., J. R. Ovenden, and R. W. G. White. 1994. Preliminary investigation of mitochondrial DNA variation in jack mackerel (Trachurus declivis, Carangidae) from south-eastern Australian waters. Marine and Freshwater Research 45:495-505.
- Jordan, A., G. Pullen, and H. Williams. 1992. Jack mackerel resource assessment in south eastern Australian waters: Final Report to FRDC, Project DFT2Z. Tasmanian Department of Primary Industry, Fisheries and Energy, Sea Fisheries Research Laboratories. Taroona, Tasmania.
- Jordan, A. R., G. Pullen, J. Marshall, and H. Williams. 1995. Temporal and spatial patterns of spawning in jack mackerel, *Trachurus declivis* (Pisces: Carangidae), during 1988-91 in eastern Tasmanian waters. Marine and Freshwater Research **46**:831-842.
- Kailola, P. J., M. J. Williams, P. C. Stewart, A. M. Reichelt, and C. Grieve. 1993. Australian Fisheries Resources. Bureau of Resource Sciences, Department of Primary Industries and Energy and the Fisheries Research and Development Corporation, Canberra.
- Keane, J. P. 2009. Mesoscale Characterisation of the Pelagic Shelf Ecosystem of South-Eastern Australia: Integrated Approach Using Larval Fish Assemblages and Oceanography. PhD Thesis, University of Tasmania. 269 pp.
- Kerstan, M. 2000. Estimation of precise ages from the marginal increment widths of differently growing sardine (*Sardinops sagax*) otoliths. Fisheries Research **46**:207-225.
- Lasker, R. 1985. An egg production method for estimating spawning biomass of pelagic fish: Application to the Northern Anchovy, *Engraulis mordax*. National Marine Fisheries Services. National Oceanic and Atmospheric Administration Technical Report.

- Last, P. R., E. O. G. Scott, and F. H. Talbot. 1983. Fishes of Tasmania. Tasmanian Fisheries Development Authority. 563 pp.
- Lindholm, R., and J. G. H. Maxwell. 1988. Stock separation of jack mackerel *Trachurus declivis* (Jenyns, 1841), and yellowtail *T. novaezealandiae* (Richardson, 1843) in southern Australian waters using principal component analysis. CSIRO Marine Laboratories Report No. 189. Hobart, Tasmania. 7 pp.
- Logerwell, E. A., B. Lavaniegos, and P. E. Smith. 2001. Spatially-explicit bioenergetics of Pacific sardine in the Southern California Bight: are mesoscale eddies areas of exceptional prerecruit production? Progress In Oceanography **49**:391-406.
- Louw, G. G., C. D. Van der Lingen, and M. J. Gibbons. 1998. Differential feeding by sardine *Sardinops sagax* and anchovy *Engraulis capensis* recruits in mixed shoals. South African Journal of Marine Science **19**:227-232.
- Lyle, J. M., K. Krusic-Golub, and A. K. Morison. 2000. Age and growth of jack mackerel and the age structure of the jack mackerel purse-seine catch. Final Report for FRDC Project 1995/034. Tasmanian Aquaculture and Fisheries Institute, University of Tasmania. Hobart, Tasmania. 49 pp.
- Markina, N. P., and V. Z. Boldyrev. 1980. Feeding of the redbait on underwater elevations of the southwest Pacific. Biologiya Morya (Vladivostok) **4**:40-45.
- Marriott, P. M., and M. J. Manning. 2011. Reviewing and refining the method for estimating blue mackerel (*Scomber australasicus*) ages. New Zealand Fisheries Assessment Report **2011**:25.
- Marshall, J. M., G. Pullen, and A. R. Jordan. 1993. Reproductive biology and sexual maturity of female jack mackerel, *Trachurus declivis* (Jenyns), in eastern Tasmanian waters. Australian Journal of Marine and Freshwater Research **44**:799-809.
- Maxwell, J. G. H. 1979. Jack mackerel. Fishery Situation Report, no. 2. Division of Fisheries and Oceanography. CSIRO Australia, 18 pp.
- May, J. L., and J. G. H. Maxwell. 1986. Field guide to trawl fish from temperate waters of Australia. CSIRO Division of Fisheries Research.
- McGarvey, R. and M. A. Kinloch. 2001. An analysis of the sensitivity of stock biomass estimates derived from the daily egg production method (DEPM) to uncertainty in egg mortality rates. Fisheries Research **49**:303-307.
- McLeod, D. J., A. J. Hobday, J. M. Lyle, and D. C. Welsford. 2012. A prey-related shift in the abundance of small pelagic fish in eastern Tasmania? ICES Journal of Marine Science: Journal du Conseil.
- Mel'nikov, Y. and N. A. Ivanin. 1995. Age-size composition and mortality of the rubyfish *Plagiogeneion rubiginosum* (Emmelichthyidae) in West Indian Submarine Ridge. Journal of Ichthyology **35**:20-27.
- Meléndez, C. R. and M. R. Céspedes. 1986. *Emmelichthys nitidus cyanescens* (Guichenot, 1848) in the Chilean Southern fisheries (Perciformes, *Emmelichthyidae*). Investigacion Pesquera **33**:111-114.
- Meyer, M. and M. J. Smale. 1991. Predation patterns of demersal teleosts from the Cape south and west coasts of South Africa. 1. Pelagic predators. South African Journal of Marine Science **10**:173-191.
- Moore, A., and M. Skirtun. 2012. The Small Pelagic Fishery. In J. Woodhams, S. Vieira and J. Stobutzki (Eds), Fishery Status Reports 2011, Australian Bureau of Agricultural and Resources Economics and Sciences, Canberra, 2012, 93 pp.
- MPI. 2022. Blue Mackerel (EMA). Fisheries Infosite. Ministry for Primary Industries. CC BY 4.0. URL: https://fs.fish.govt.nz/Page.aspx?pk=7&tk=100&sc=EMA. Accessed 10 May 2022.

- Murphy, J. J., F. A. Ochwada-Doyle, L. D. West, K. E. Stark, and J. M. Hughes. 2020. The NSW Recreational Fisheries Monitoring Program - survey of recreational fishing, 2017/18. NSW DPI - Fisheries Final Report Series No. 158.
- Neira, F. J. 2011. Application of daily egg production to estimate biomass of jack mackerel, *Trachurus declivis* - a key fish species in the pelagic ecosystem of south-eastern Australia. Final Report to the Winifred Violet Scott Charitable Trust. Fisheries, Aquaculture and Coasts Centre, Institute for Marine and Antarctic Studies (IMAS), University of Tasmania. 42 pp.
- Neira, F. J., J. M. Lyle, G. P. Ewing, J. P. Keane, and S. R. Tracey. 2008. Evaluation of egg production as a method of estimating spawning biomass of redbait off the east coast of Tasmania. Final report, FRDC project no. 2004/039. Tasmanian Aquaculture and Fisheries Institute, Hobart.
- Neira, F. J., A. Miskiewicz, and T. Trnski. 1998. Larvae of temperate Australian fishes: laboratory guide for larval fish identification. University of Western Australia Press, Nedlands, W.A.
- Neira, F. J., M. I. Sporcic, and A. R. Longmore. 1999. Biology and fishery of pilchard, *Sardinops sagax* (Clupeidae), within a large south-eastern Australian bay. Marine and Freshwater Research **50**:43-55.
- Nelson, J. S. 2006. Fishes of the World. Fourth Edition. John Wiley and Sons, Inc. Hoboken, New Jersey. 601 pp.
- Parin, N. V., A. N. Mironov, and K. N. Nesis. 1997. Biology of the Nazca and Sala y Gomez submarine ridges, an outpost of the Indo-West Pacific Fauna in the Eastern Pacific Ocean: composition and distribution of the fauna, its communities and history. Advances in Marine Biology **32**:145-242.
- Parker, K. 1980. A direct method for estimating northern anchovy, *Engraulis mordax*, spawning biomass. Fisheries Bulletin **84**:541-544.
- Parrish, R. H., R. Serra, and W. S. Grant. 1989. The Monotypic Sardines, Sardina and Sardinops -Their Taxonomy, Distribution, Stock Structure, and Zoogeography. Canadian Journal of Fisheries and Aquatic Sciences 46:2019-2036.
- Patterson, H., D. Bromhead, D. Galeano, J. Larcombe, J. Woodhams, and R. Curtotti. 2021. Fishery status reports 2021. Australian Bureau of Agricultural and Resource Economics and Sciences, Canberra. CC BY 4.0.
- Paul, L. J., P. L. Horn, and M. P. Francis. 2000. Development of an ageing methodology, and first estimates of growth parameters and natural mortality for rubyfish (*Plagiogeneion rubiginosum*) off the east coast of the North Island (QMA 2). Fisheries Assessment Report 2000/22. Ministry of Fisheries, New Zealand.
- Piddocke, T., C. Ashby, K. Hartmann, A. Hesp, P. Hone, J. Klemke, S. Mayfield, A. Roelofs, T. Saunders, J. Stewart, B. Wise, and J. Woodhams (eds). 2021. Status of Australian fish stocks reports 2020, Fisheries Research and Development Corporation, Canberra.
- Prokopchuk, I., and E. Sentyabov. 2006. Diets of herring, mackerel, and blue whiting in the Norwegian Sea in relation to *Calanus finmarchicus* distribution and temperature conditions. ICES Journal of Marine Science **63**:117-127.
- Pullen, G. 1994. Fishery status report: Purse-seine (The Tasmanian jack mackerel fishery). Department of Primary Industry and Fisheries, Tasmania. Internal Report 13. 49pp.
- Richardson, B. 1982. Geographical distribution of electrophoretically detected protein variation in Australian commercial fishes. I. Jack mackeral, *Trachurus declivis* Jenyns. Marine and Freshwater Research **33**:917-926.
- Rogers, P. J., and T. M. Ward. 2007. Application of a 'case-building approach' to investigate the age distributions and growth dynamics of Australian sardine (*Sardinops sagax*) off South Australia. Marine and Freshwater Research **58**:461-474.

- Rogers, P. J., T. M. Ward, L. J. McLeay, M. Lowry, and D. Williams. 2009. Reproductive biology of blue mackerel, *Scomber australasicus*, off southern and eastern Australia: suitability of the Daily Egg Production Method for stock assessment. Marine and Freshwater Research 60:187-202.
- Rohde, K. 1987. Different populations of *Scomber australasicus* in New Zealand and south-eastern Australia, demonstrated by a simple method using monogenean sclerites. Journal of Fish Biology **30**:651-657.
- Schmarr, D. W., I. D. Whittington, J. R. Ovenden, and T. M. Ward. 2011. Discriminating stocks of Scomber australasicus using a holistic approach: a pilot study. American Fisheries Society Symposium 76:1-21.
- Schwartzlose, R. A., J. Alheit, A. Bakun, T. R. Baumgartner, R. Cloete, R. J. M. Crawford, W. J. Fletcher, Y. Green-Ruiz, E. Hagen, T. Kawasaki, D. Lluch-Belda, S. E. Lluch-Cota, A. D. MacCall, Y. Matsuura, M. O. Nevárez-Martínez, R. H. Parrish, C. Roy, R. Serra, K. V. Shust, M. N. Ward, and J. Z. Zuzunaga. 1999. Worldwide large-scale fluctuations of sardine and anchovy populations. South African Journal of Marine Science 21:289-347.
- Scoles, D. R., B. B. Collette, and J. E. Graves. 1998. Global phylogeography of mackerels of the genus *Scomber*. Fishery Bulletin **96**:823-842.
- Sexton, S. C., T. M. Ward, and C. Huveneers. 2017. Characterising the spawning patterns of Jack Mackerel (*Trachurus declivis*) off eastern Australia to optimise future survey design. Fisheries Research **186**:223-236.
- Shuntov, V. P. 1969. Some features of the ecology of pelagic fishes in the Great Australian Bight. Problems of Ichthyology **9**:801-809.
- Smith, A. D. M., T. M. Ward, F. Hurtado, N. Klaer, E. Fulton, and A. E. Punt. 2015. Review and update of harvest strategy settings for the Commonwealth Small Pelagic Fishery: Single species and ecosystem considerations. Final Report of FRDC Project No. 2013/028. CSIRO Oceans and Atmosphere Flagship, Hobart. 74 pp.
- Smith, K. A., and K. Deguara. 2003. Formation and annual periodicity of opaque zones in sagittal otoliths of *Mugil cephalus* (Pisces: Mugilidae). Marine and Freshwater Research **54**:57-67.
- Smith, M. M., and P. C. Heemstra. 1986. Smith's Sea Fishes. MacMillan, Johannesburg. 1047 pp.
- Smolenski, A. J., J. R. Ovenden, and R. W. G. White. 1994. Preliminary investigation of mitochondrial DNA variation in jack mackerel (*Trachurus declivis*, Carangidae) from south-eastern Australian waters. Marine and Freshwater Research 45:495-505.
- Stevens, J. D., and H. F. Hausfeld. 1982. Age determination and mortality estimates of an unexploited population of Jack mackerel *Trachurus declivis* (Jenyns, 1841) from south-east Australia. CSIRO Marine Laboratories Report. Hobart, Tasmania. 148 pp.
- Stevens, J. D., H. F. Hausfeld, and S. R. Davenport. 1984. Observations on the biology, distribution and abundance of *Trachurus declivis*, *Sardinops neopilchardus* and *Scomber australasicus* in the Great Australian Bight. CSIRO Marine Laboratories, Cronulla.
- Stewart, J., and D. J. Ferrell. 2001. Age, growth, and commercial landings of yellowtail scad (*Trachurus novaezelandiae*) and blue mackerel (*Scomber australasicus*) off the coast of New South Wales, Australia. New Zealand Journal of Marine and Freshwater Research **35**:541-551.
- Stewart, J., D. J. Ferrell, and N. L. Andrew. 1999. Validation of the formation and appearance of annual marks in the otoliths of yellowtail (*Trachurus novaezelandiae*) and blue mackerel (*Scomber australasicus*) in New South Wales. Marine and Freshwater Research **50**:389-395.
- Stratoudakis, Y., M. Bernal, K. Ganias, and A. Uriarte. 2006. The daily egg production method: recent advances, current applications and future challenges. Fish and Fisheries **7**:35-57.

- Tracey, S., C. Buxton, C. Gardner, B. Green, K. Hartmann, M. Haward, J. Jabour, J. Lyle, and J. McDonald. 2013. Super Trawler Scuppered in Australian Fisheries Management Reform. Fisheries 38:345-350.
- van der Lingen, C. D. 1994. Effect of Particle-Size and Concentration on the Feeding-Behavior of Adult Pilchard *Sardinops sagax*. Marine Ecology-Progress Series **109**:1-13.
- van der Lingen, C. D., and J. A. Huggett. 2003. The role of ichthyoplankton surveys in recruitment research and management of South African anchovy and sardine.*in* H. I. Browman and A. B. Skiftesvic, editors. The Big Fish Bang. Proceedings of the 26th Annual Larval Fish Conference. Institute of Marine Research, Norway.
- Waldron, M. 1998. Annual ring validation of the South African sardine *Sardinops sagax* using daily growth increments. South African Journal of Marine Science **19**:425-430.
- Ward, P. J., T. Timmiss, and B. Wise. 2001a. A review of biology and fisheries for mackerel. Bureau of Rural Sciences, Canberra.
- Ward, T., B. Molony, J. Stewart , J. Andrews, and A. Moore. 2014a. Australian Sardine Sardinops sagax, in M. Flood, I. Stobutzki, J. Andrews, C. Ashby, G. Begg, R. Fletcher, C. Gardner, L. Georgeson, S. Hansen, K. Hartmann, P. Hone, P. Horvat, L. Maloney, B. McDonald, A. Moore, A. Roelofs, K. Sainsbury, T. Saunders, T. Smith, C. Stewardson, .J Stewart & B. Wise (eds) 2014, Status of key Australian fish stocks reports 2014, Fisheries Research and Development Corporation, Canberra.
- Ward, T. M., P. Burch, L. J. McLeay, and A. R. Ivey. 2011a. Use of the Daily Egg Production Method for stock assessment of sardine, *Sardinops sagax*; lessons learned over a decade of application off southern Australia Reviews in Fisheries Science **19**:1-20.
- Ward, T. M., O. Burnell, A. Ivey, J. Carroll, J. Keane, J. Lyle, and S. Sexton. 2015a. Summer spawning patterns and preliminary daily egg production method survey of Jack Mackerel and Australian Sardine off the East Coast. South Australian Research and Development Institute (Aquatic Sciences), Adelaide.
- Ward, T. M., O. W. Burnell, A. Ivey, S. C. Sexton, J. Carroll, J. Keane, and J. M. Lyle. 2016. Spawning biomass of jack mackerel (*Trachurus declivis*) off eastern Australia: Critical knowledge for managing a controversial fishery. Fisheries Research **179**:10-22.
- Ward, T. M. and G. L. Grammer. 2016. Commonwealth Small Pelagic Fishery: Fishery Assessment Report 2015. Report to the Australian Fisheries Management Authority South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2010/000270-7. SARDI Research Report Series No. 900. 111 pp.
- Ward, T. M. and G. L. Grammer. 2017. Commonwealth Small Pelagic Fishery: Fishery Assessment Report 2014-2016. Report to the Australian Fisheries Management Authority. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2010/000270-8. SARDI Research Report Series No. 948. 123 pp.
- Ward, T. M. and G. L. Grammer. 2018. Commonwealth Small Pelagic Fishery: Fishery Assessment Report 2017. Report to the Australian Fisheries Management Authority South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2010/000270-9. SARDI Research Report Series No. 982. 114pp.
- Ward, T. M. and G. L. Grammer. 2019. Commonwealth Small Pelagic Fishery: Fishery Assessment Report 2018. Report to the Australian Fisheries Management Authority., South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2010/000270-10. SARDI Research Report Series No. 1021. 132 pp.
- Ward, T. M. and G. L. Grammer. 2021a. Commonwealth Small Pelagic Fishery: Fishery Assessment Report 2020. Report to the Australian Fisheries Management Authority., South Australian

Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2020/000209-2. SARDI Research Report Series No. 1085. 8 pp.

- Ward, T. M. and G. L. Grammer. 2021b. Commonwealth Small Pelagic Fishery: Status Summary Report 2019. Report to the Australian Fisheries Management Authority., South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2020/000209-1. SARDI Research Report Series No. 1057. 7pp.
- Ward, T. M., G. L. Grammer, and A. R. Ivey. 2021a. Spawning biomass of Blue Mackerel (*Scomber australasicus*) and Australian Sardine (*Sardinops sagax*) in the East sub-area of the Small Pelagic Fishery. Report to the Australian Fisheries Management Authority., South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2021/000047-1. SARDI Research Report Series No 1087. 54pp.
- Ward, T. M., G. L. Grammer, A. R. Ivey, J. R. Carroll, J. P. Keane, J. Stewart, and L. Litherland. 2015b. Egg distribution, reproductive parameters and spawning biomass of Blue Mackerel, Australian Sardine and Tailor off the East Coast during late winter and early spring. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. FRDC Project No. 2014/033. 77 p.
- Ward, T. M., G. L. Grammer, A. R. Ivey, and J. Keane. 2020a. Spawning biomass of Jack Mackerel (Trachurus declivis) in the East sub-area of the Small Pelagic Fishery during summer 2019. Report to the Australian Fisheries Management Authority. . South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2020/000206-01. 42pp.
- Ward, T. M., G. L. Grammer, A. R. Ivey, and J. P. Keane. 2019. Spawning biomass of Redbait (*Emmelichthys nitidus*) between western Kangaroo Island, South Australia and southwestern Tasmania in October 2017. Report to the Australian Fisheries Management Authority., South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2019/000053-1. SARDI Research Report Series No. 1011. 38pp.
- Ward, T. M., G. L. Grammer, A. R. Ivey, J. J. Smart, and J. P. Keane. 2018. Spawning biomass of Jack Mackerel (*Trachurus declivis*) and Sardine (*Sardinops sagax*) between western Kangaroo Island, South Australia and south-western Tasmania. Report to the Australian Fisheries Management Authority. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2018/000174-1. SARDI Research Report Series No. 983. 51pp.
- Ward, T. M., G. L. Grammer, A. R. Ivey, J. J. Smart, and R. McGarvey. 2021b. Increasing the precision of the daily egg production method; 2020's remix of a 1980's classic. ICES Journal of Marine Science **78**:1177-1195.
- Ward, T. M., F. Hoedt, L. McLeay, W. F. Dimmlich, M. Kinloch, G. Jackson, R. McGarvey, P. J. Rogers, and K. Jones. 2001b. Effects of the 1995 and 1998 mass mortality events on the spawning biomass of *Sardinops sagax* in South Australian waters. ICES Journal of Marine Science **58**:830-841.
- Ward, T. M., A. R. Ivey, and J. Earl. 2014b. Commonwealth Small Pelagic Fishery: Fishery Assessment Report 2013. Report to the Australian Fisheries Management Authority., South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2010/000270-5. SARDI Research Report Series No. 778. 105 pp.
- Ward, T. M., A. R. Ivey, and J. Earl. 2015c. Commonwealth Small Pelagic Fishery: Fishery Assessment Report 2014. Report to the Australian Fisheries Management Authority. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2010/000270-6. SARDI Research Report Series No. 847. 113 pp.

- Ward, T. M., A. R. Ivey, and D. Gorman. 2013. Commonwealth Small Pelagic Fishery: Fishery Assessment Report 2012. Report to the Australian Fisheries Management Authority. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2010/000270-4. SARDI Research Report Series No. 696. 100 pp.
- Ward, T. M., J. Lyle, J. P. Keane, G. A. Begg, P. Hobsbawn, A. R. Ivey, R. Sakabe, and M. A. Steer. 2011b. Commonwealth Small Pelagic Fishery: Status Report 2010. Report to the Australian Fisheries Management Authority. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2010/000270-2. SARDI Research Report Series No. 524. 84 pp.
- Ward, T. M., J. Lyle, J. P. Keane, G. A. Begg, P. Hobsbawn, A. R. Ivey, R. Sakabe, and M. A. Steer.
  2012. Commonwealth Small Pelagic Fishery: Fishery Assessment Report 2011. Report to the Australian Fisheries Management Authority. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2010/000270-3. SARDI Research Report Series No. 614. 98 pp.
- Ward, T. M., R. McGarvey, A. Ivey, and G. L. Grammer. 2020b. Validating a new sampling technique for estimating egg production. South Australian Research and Development institute (SARDI), Adelaide. FRDC Project 2017-027. 33pp. .
- Ward, T. M., L. J. McLeay, P. J. Rogers, and W. F. Dimmlich. 2001c. Spawning Biomass of Pilchard (*Sardinops sagax*) in South Australia in 2001. Report to Primary Industries and Resourses South Australia Fisheries. South Australian Research and Development Institute (Aquatic Sciences).
- Ward, T. M., and P. J. Rogers, editors. 2007. Development and evaluation of egg-based stock assessment methods for blue mackerel *Scomber australasicus* in southern Australia. Final report to the Fisheries Research and Development Corporation Project 2002/061. SARDI Aquatic Science, Adelaide.
- Ward, T. M., P. J. Rogers, L. J. McLeay, and R. McGarvey. 2009. Evaluating the use of the daily egg production method for stock assessment of blue mackerel, *Scomber australasicus*. Journal of Marine and Freshwater Research **62**:112-128.
- Ward, T. M., D. W. Schmarr, L. J. McLeay, P. J. Rogers, and A. Ivey. 2007. A preliminary investigation of the spawning biomass of sardine (pilchard, *Sardinops sagax*) off eastern Australia. Final report for the New South Wales Department of Primary Industry. South Australian Research and Development Institute (Aquatic Sciences), Adelaide.
- Ward, T. M., J. Smart, G. L. Grammer, A. Ivey, and R. McGarvey. 2020c. Stock assessment of Australian Sardine (*Sardinops sagax*) off South Australia 2019. Report to PIRSA Fisheries and Aquaculture. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2007/000765-7. SARDI Research Report Series No. 1048. 107 pp.
- Ward, T. M., and J. Staunton-Smith. 2002. Comparison of the spawning patterns and fisheries biology of the sardine, *Sardinops sagax*, in temperate South Australia and sub-tropical southern Queensland. Fisheries Research **56**:37-49.
- Watanabe, Y., H. Zenitani, and R. Kimura. 1996. Offshore expansion of spawning of the Japanese sardine, *Sardinops melanostictus*, and its implication for egg and larval survival. Canadian Journal of Fisheries and Aquatic Sciences **53**:55-61.
- Webb, B. F. 1976. Aspects of the biology of jack mackerel, *Trachurus declivis* (Jenyns) from southeast Australian waters. Tasmanian Fisheries Research **10**:1-14.
- Welsford, D. C., and J. M. Lyle. 2003. Redbait (*Emmelichthys nitidus*): A synopsis of fishery and biological data. TAFI Technical Report Series, no. 20 20

- Whitehead, P. J. P. 1985. FAO Species Catalogue. Vol. 7. Clupeoid fishes of the world (suborder Clupeoidei). An annotated and illustrated catalogue of the herrings, sardines, pilchards, sprats, shads, anchovies and wolf-herrings. FAO Fish. Synop. 125(7/1):1-303. Rome: FAO.
- Whittington, I. D., M. Crockford, D. Jordan, and B. Jones. 2008. Herpesvirus that caused epizootic mortality in 1995 and 1998 in pilchard, *Sardinops sagax* (Steindachner), in Australia is now endemic. Journal of Fish Diseases **31**:97-105.
- Williams, H., and G. Pullen. 1986. A synopsis of biological data on the jack mackerel *Trachurus declivis* Jenyns. Technical Report No. 10. Department of Sea Fisheries, Tasmania, Marine Laboratories: Taroona. 34 pp.
- Young, J. W., A. R. Jordan, C. Bobbi, R. E. Johannes, K. Haskard, and G. Pullen. 1993. Seasonal and interannual variability in krill (*Nyctiphanes australis*) stocks and their relationship to the fishery for jack mackerel (*Trachurus declivis*) off eastern Tasmania, Australia. Marine Biology 116:9-18.